

## EFFECT OF CAMPYLOBACTER COLI LPS ON PHAGOCYTOSIS IN VITRO AND IN VIVO

Dunya Fareed<sup>\*1</sup>, Rasmia Abed Abu-Risha<sup>1</sup> and Ali Hafeedh Abbas<sup>2</sup>

<sup>1</sup>Biology Department, College of Sciences, University of Baghdad, Baghdad-Iraq

<sup>2</sup>University of Baghdad – College of Sciences – Tropical Biological Researches Unit.

Article Received on  
05 April 2014,

Revised on 28 April 2014,  
Accepted on 21 May 2014

**\*Correspondence for  
Author**

**Dr. Dunya Fareed**

Biology Department, College  
of Sciences, University of  
Baghdad, Baghdad-Iraq

### ABSTRACT

*Campylobacter coli* usually live in the intestinal tract of animals and cause inflammation of the intestine and diarrhea. In this study, *C. coli* were isolated from patients having diarrhea and identified based on morphological and biochemical characteristics. Lipopolysaccharide (LPS) is major outer membrane structural component of gram negative bacteria which induces a broad range of biological response. Here, LPS was isolated with hot phenol-water extraction and purified by gel filtration using Sephadex G-200. Phagocytosis is essential for fighting infections in immune responses to protect organisms from bacteria invasion hence we studied the effect of purified LPS on phagocytosis

both *in vitro* and *in vivo*. The results reported in this study may prove to be valuable in analyzing the LPS effect on host immune response.

**KEYWORDS:** *C. coli*, LPS, Phagocytosis.

### INTRODUCTION

Campylobacter species are among the most commonly reported bacterial cause of acute gastroenteritis disease worldwide and especially in the industrialized countries.<sup>[1, 2]</sup> Approximately nine million human campylobacteriosis cases are reported annually in 27 European countries.<sup>[3]</sup> Although most of the human campylobacteriosis is caused by *C. jejuni*, the importance of *C. coli* study is being recognized due to an increasing resistance of this species to number of antimicrobials.<sup>[4]</sup> Most Campylobacter research focuses upon *C. jejuni* but still campylobacteriosis caused by *C. coli* infection results in millions dollars of annual cost.<sup>[5, 6]</sup>

Livestock wastes from intensive animal production such as manure or slurry contain pathogenic microorganisms including *Campylobacter* species.<sup>[7]</sup> It has been reported that drinking water may be the source of *C. coli* infection in grandparent breeder farms.<sup>[8]</sup> This has increased concern about effect of pathogens in animal manure on human and animal health.<sup>[9]</sup> For *Campylobacter* motility is important for intestinal colonization and invasion of epithelial cells. In *Campylobacter* species flagella function as secretory organelles and recognized as a major virulence factor for this pathogen.<sup>[10, 11]</sup> It is reported earlier that the flagellar structural protein (FlaA) is immunoprotective antigen,<sup>[12, 13]</sup> flagellins are heavily glycosylated proteins and are highly immunogenic.<sup>[14, 15]</sup>

Macrophages play an important role in recognition, uptake and killing of parasites during pathogen infections.<sup>[16]</sup> It has been reported that LPS activates several signal transduction pathways in macrophages leading to various gene activation and cytokine production.<sup>[17, 18]</sup> In this study we have isolated LPS from *C. coli* and detected its effect on Phagocytosis using human macrophages: both *in vivo* and *in vitro*.

## MATERIALS AND METHODS

### Isolation of *C. coli*

*C. coli* were isolated from diarrhea samples collected from Al-Yarmok Hospital Baghdad, Iraq. The serially diluted samples were inoculated on blood agar containing 5 % SRBCs and other supplement and incubated for 48 hours at 4 °C under micro aerobic conditions. The single isolated colonies were identified by morphological characteristics, gram Staining and biochemical tests.

### Extraction of LPS from *C. coli*

The LPS was extracted from *C. coli* as described previously by Preston and Penner.<sup>[19]</sup> Briefly digestive enzymes were used to isolate LPS and hot phenol method was used for extraction process. The extracted LPS were then purified by gel filtration chromatography using Sephadex G-200 as described by manufacturer.

### Blood sample preparation

Two milliliter of blood samples was collected from twenty healthy persons in heparinized (50 IU/ml) silicon test tube to prevent the absorption of cells on glass tubes (according to WHO guideline). These Blood samples were directly used to evaluate phagocytosis.

### *In vitro* phagocytosis assay

The *in vitro* phagocytosis was determined by procedure described by Sun<sup>[20]</sup> Briefly, monocytes were added into the wells and after 6 h of incubation the floating cells were removed and monocytes were washed twice with PBS. Under the phase microscope, about 300 macrophages per sample including LPS engulfed were counted and the percentage ratio was calculated as the relative number of cells that engulfed LPS per total cells.

#### ***In vivo* phagocytosis assay**

The *in vivo* phagocytosis was estimated as described by Tanaka.<sup>[21]</sup> Briefly LPS was diluted 10-fold with PBS and injected into the mouse peritoneal cavity by 10  $\mu$ l/g body weight. After overnight starvation the mice were sacrificed and macrophages were recovered from Peritoneal, Liver and Spleen. For each sample 400 cells were counted, and the percentage ratio was calculated as the relative number of cells that engulfed LPS per total cells.


### **RESULTS AND DISCUSSION**

*C. coli* were isolated from patients having diarrhea. The morphological characteristics such as colony morphology and motility were observed and gram staining with biochemical tests such as oxidase and hiturate were performed for identification of *C. coli*. The observed results were summarized in Table 1.

**Table 1: Biochemical test and characterization of *C. coli***


Test	Result
Colony on blood agar	Appear small convert grey white colony
Gram staining	Negative (Gram negative bacteria)
Oxidase test	Positive
Hiturate test	Hydrolysis negative
Motility	Motile
Shape	Spiral shape

LPS are heat stable endotoxins and long been known to induce variety of pathophysiologic responses, including fever, coagulant activity, septic shock, and death.<sup>[22]</sup> Here, we have used hot phenol method to extract LPS from the isolated *C. coli*. To remove the impurities such as protein and RNA, extracted samples were further purified by Size exclusion chromatography using Sephadex G 200. After purification absorbance was measured at 490 nm for collected fractions (Figure 1). The fraction showing maximum absorbance was used for further phagocytosis study.




**Figure 1: Purification of LPS by gel filtration (Sephadex G-200)**

Phagocytosis is controlled by a complex signals located on the target cell surface and their respective receptors which determine phagocytosis process. [23] Many approaches have been designed to study these processes using macrophages as prototypic inflammatory cells, since these cells are potently activated by LPS. [24] In present investigation we studied phagocytosis both *in vivo* and *in vitro*. The percentage ratio was determined for phagocytosed LPS and presented in figure 2 for *in vitro* study and in figure 3 for *in vivo* study.



**Figure 2: Phagocytosis *in vitro***



**Figure 3: Phagocytosis *in vivo***

In presence of LPS increased phagocytosis was observed in both *in vitro* and *in vivo* studies as compared to control which indicates that LPS exposure required for phagocytosis progress. The increased phagocytosis reported in this study may prove to be valuable in analyzing the alterations induced by LPS on macrophage involved in the host response.

## REFERENCES

1. Friedman CR, Neimann J, Wegener HC, Tauxe R. Epidemiology of *Campylobacter jejuni* infection in the United States and other industrialized nations, In: *Campylobacter* (eds) I Nachamkin I and MJ Blaser (Washington: ASM Press): 2001.
2. World Health Organization (WHO). The increasing incidence of human campylobacteriosis. Copenhagen, Denmark. 2000, pp. 21-25.
3. Andreoletti O, Budka H, Buncic S, Collins JD, Griffin J, Hald T *et al.* Scientific opinion on *Campylobacter* in broiler meat production: control options and performance objectives and/or targets at different stages of the food chain. *EFSA J*, 2011; 9: 141.
4. Gebreyes WA, Thakur S, Morrow WEM. *Campylobacter coli*: prevalence and antimicrobial resistance in antimicrobial-free (ABF) swine production systems. *J. Antimicrob Chemother*, 2005; 56: 765–8.
5. Humphrey T, O'Brien S, Madsen M. *Campylobacter* as zoonotic pathogens: a food production perspective. *Int. J. Food Microbiol*, 2007; 117: 237–57.
6. Sheppard SK, Dallas JF, Wilson DJ, Strachan NJC, McCarthy ND, Jolley KA *et al.* Evolution of an agriculture associated disease causing *Campylobacter coli* clade:

- evidence from national surveillance data in Scotland. *PLoS ONE*, 2010; 5: e15708 doi:10.1371/journal.pone.0015708.
7. Klein M, Brown L, Ashbolt NJ, Stuetz RM, Roser DJ. Inactivation of indicators and pathogens in cattle feedlot manures and compost as determined by molecular and culture assays. *FEMS Microbiol Ecol*, 2011; 77: 200–10.
  8. Pérez-Boto D, García-Peña FJ, AbadMoreno JC, Hurtado-Pizarro MD, Pérez-Cobo I, Aurora Echeita M. Drinking water as the source of *Campylobacter coli* infection in grandparent heavy breeders. *Avian Pathol*, 2010; 39: 483–7.
  9. Bicudo JR, Goyal SM. Pathogens and manure management systems: a review. *Environ. Technol*, 2003; 24: 115–30.
  10. Konkel ME, Klena JD, Rivera-Amill V, Monteville MR, Biswas D, Raphael B, Mickelson J. Secretion of virulence proteins from *Campylobacter jejuni* is dependent on a functional flagellar export apparatus. *J Bacteriol*, 2004; 186: 3296–303.
  11. Song YC, Jin S, Louie H, Ng D, Lau R, Zhang Y. FlaC, a protein of *Campylobacter jejuni* TGH9011 (ATCC43431) secreted through the flagellar apparatus, binds epithelial cells and influences cell invasion. *Mol Microbiol*, 2004; 53: 541–53.
  12. Guerry P, Alm RA, Power ME, Logan SM, Trust TJ. Role of two flagellin genes in *Campylobacter* motility. *J Bacteriol*, 1991; 173: 4757–64.
  13. Martin PM, Mathiot J, Ipero J, Kirimath M, Georges AJ, Georges-Courbot MC. Immune response to *Campylobacter jejuni* and *Campylobacter coli* in a cohort of children from birth to 2 years of age. *Infect Immun*, 1989; 57: 2542–6.
  14. Logan SM, Kelly JF, Thibault P, Ewing CP, Guerry P. Structural heterogeneity of carbohydrate modifications affects serospecificity of *Campylobacter* flagellins. *Mol Microbiol*, 2002; 46: 587–97.
  15. Thibault P, Logan SM, Kelly JF, Brisson JR, Ewing CP, Trust TJ, Guerry P. Identification of the carbohydrate moieties and glycosylation motifs in *Campylobacter jejuni* flagellin. *J Biol Chem*, 2001; 276: 34862–70.
  16. Hoebe K, Janssen B, Beutler B. The interface between innate and adaptive immunity. *Nat Immunol*, 2004; 5: 971–4.
  17. Raetz CR, Ulevitch RJ, Wright SD, Sibley CH, Ding A, Nathan CF. Gram-negative endotoxin: an extraordinary lipid with profound effects on eukaryotic signal transduction. *FASEB J*, 1991; 5: 2652–60.
  18. Sweet MJ, Hume DA. Endotoxin signal transduction in macrophages. *J Leukocyte Biol*, 1996; 60: 8–26.

19. Preston MA, Penner JL. Structural and antigenic properties of lipopolysaccharides from serotype reference strains of *Campylobacter jejuni*. *Infection and immunity*, 1987; 55: 1806-12.
20. Sun SF, Pan QZ, Hui X, Zhang BL, Wu HM, Li H. Stronger in vitro phagocytosis by monocytes-macrophages is indicative of greater pathogen clearance and antibody levels in vivo. *Poult Sci*, 2008; 87: 1725-33.
21. Tanaka N, Abe-Dohmae S, Iwamoto N, Fitzgerald ML, Yokoyama S. HMG-CoA reductase inhibitors enhance phagocytosis by upregulating ATP-binding cassette transporter A7. *Atherosclerosis*, 2011; 217: 407-14.
22. Schletter J, Heine H, Ulmer AJ, Rietschel ET. Molecular mechanisms of endotoxin activity. *Arch Microbiol*, 1995; 164: 383-9.
23. Ravichandran KS. Beginnings of a good apoptotic meal: the find-me and eat-me signaling pathways. *Immunity*, 2011; 35: 445–55.
24. Morrison DC, Danner RL, Dinarello CA, Munford RS, Natanson C, Pollack M *et al*. Bacterial endotoxins and pathogenesis of Gram-negative infections: current status and future directions. *J Endotoxin Res*, 1994; 1: 71-83.