

**PHYSICOCHEMICAL ANALYSIS OF PT.ET. EXTRACTED
LYCOPENE AND VISCOSITY, SURFACE TENSION
CHARACTERIZATION BY MANSINGH SURVISMETER**

Anil Kumar¹, Bijendra Singh² and Kapil Tyagi³

¹Centre for Nanosciences, Central University of Gujarat, Gandhinagar, Gujarat, India.

²School of Chemical Sciences, Central University of Gujarat, Gandhinagar, Gujarat, India.

³Department of Chemistry, Mewar University, Chittorgarh, Rajasthan, India

Article Received on
05 April 2014,

Revised on 28 April 2014,
Accepted on 21 May 2014

***Correspondence for
Author**

Anil Kumar

Centre for Nanosciences,
Central University of Gujarat,
Gandhinagar, Gujarat, India.

ABSTRACT

The Physico-chemical characterization of Lycopene by applied analysis methods such as Absorbance, pH, Viscosity and Surface Tension, Lycopene is a lipid molecules substrate present in vegetables fruits for responsible to red color appeared with nutritional and medicinal properties, mostly used in skin cancer and Lycopene is protective in several chronic diseases and also Lycopene have phytochemical properties, that's way determined to applied parameter values. Quantities of Lycopene in samples CS1, AS2 to 0.042783 and 0.016227 in mg/100gm, viscosity of CS1, AS2 to 0.400467 N.sm⁻² and 0.371466 N.sm⁻² and surface tension of CS1, AS2 to 20.59763N.m⁻¹

and 22.87903 N.m⁻¹ and the pH values was CS1 of 6.25, AS2 of 6.60 and pure PtEt of 6.77.

Keywords: PtEt, Lycopene, pH, Absorbance, Viscosity, Surface Tension.

INTRODUCTION

The Lycopene has looked at its effect on cancer, there is growing evidence that Lycopene is protective in several chronic diseases including cardiovascular disease [1-2]. Lycopene belongs to the family of carotenoids. It has a structure that consists of a long chain of conjugated double bonds, with two open end rings. The structure Lycopene is the longest of all carotenoids. The Lycopene ([C₄₀H₅₆], MW 536.85) is an unsaturated hydrocarbon carotenoid containing 13 C=C bonds, 11 of which are conjugated and arranged in a linear array. These conjugated double bonds are responsible for the vibrant red color of Lycopene [3]. Lycopene the red pigment of the tomato is a C₄₀ carotenoid made up of eight Isoprene

unit making it a tetraterpene β - carotene the yellow pigment of the carrot is an isomer of Lycopene in which the double bond at C_1-C_2 and $*C_1- *C_2$ are replaced by bond extending from C_1 to C_6 and from $*C_1$ to $*C_6$ from rings, and also constituent of the tomatoes' This is the first step in the overall visual cycle associated with night vision. Lycopene is a vibrant red carotenoid that serves as an intermediate for the biosynthesis of other carotenoids and is found in moderate to high concentrations in such foods as tomato, watermelon, red grapefruit, and Brazilian guava etc [4]. The major coloring principle of Lycopene extract from tomato is all-*trans*-lycopene, products consists predominantly of all-*trans*-lycopene 35-96% of the total lycopene content and low levels of *cis*-lycopenes 1-22% of the total lycopene content [5]. Like its biosynthetic derivatives such as β -carotene, Lycopene free radical scavenger [6] and its presence in the diet positively correlates with reduced cancer incidence [7-10]. Conventional spectrometer assays employ a release of Lycopene from tissue with polar organic solvents followed by extraction of released Lycopene into a nonpolar solvent [11]. In a fallow up lab will examine the UV-Vis.spectrum of Lycopene isomerizes it and then examine the isomerize spectrum for comparison [12]. Lycopene extract from tomato is a dark-red viscous liquid. It is freely soluble in acetone and PtEt (Petroleum Ether) and partially soluble in ethanol and acetone, and insoluble in water. Spectroscopy was used for quantification of Lycopene in tomato varieties [13-14].

Research Methodology [15].

Collection of Sample: Two types the tomatoes were purchased from a store so variety name were not available of the tomatoes used look like cherry (CS1) and aroma (AS2) types.

Sample Preparation: 100gm of the tomatoes fruit tissue was cut in to small pieces approximately 2-3cm cubes and homogenized with Acetone using mortar and pestle until the residue is color less. Pool the Acetone extract and transfer in a separating funnel containing about 20ml PtEt and mix gently. Add about 20ml of 5% $NaSO_4$ solution and shake the separating funnel gently. PtEt to the separating funnel for clear separation of two phases most of the colors will be noticed in appear PtEt phase and re-extract the lower aqueous phases with additional 20ml PtEt until the aqueous phases is colorless. That's way the total volume of extracted lycopene at 40ml its storage in freeze for long time to use.

pH Values of Extracted Lycopene: the pH values played an important role determine the potential numbers of hydrogen of anhydrous liquid samples. The samples of PtEt extracted lycopene know pH values at 20°C by using pH meter.




Absorbance of Extracted Lycopene: Make up the volume and measure the Absorbance by spectrophotometer at 503nm (λ) using PtEt as blank [16] The carotenoid in the sample are extracted in Acetone and then taken up in the PtEt lycopene has absorption maxima at 503nm (λ) one mole at lycopene when dissolve in one little light PtEt 40-60°C and measured in a spectrophotometer at 503nm (λ) to 1cm light path gives on absorbance of $17.2 \times 10^4 / M \times \text{cm}$. There for a concentration of 3.1206 μg lycopene/ml or Lycopene in mg/100gm = $A_{503} * 3.1206 / \text{wt of sample}$ [17-18]


Determine the Viscosity and surface tension of Extracted Lycopene by BMS: [19]

The survismeter is an instrument for determining viscosity, surface tension and other parameters more than twenty five may be characterizing of various types of liquids. First maintained temperate of systems at 15-20°C and filled 30ml PtEt extracted Lycopene is in bulb of viscosity meter up to the given mark with the help of syringe after that note the time of viscosity of PtEt through the stopwatch after that know surface tension 30ml PtEt extracted Lycopene samples was filled in bulb of Stalagmometer up to the given mark with the help of syringe after that note the PDN details showing in figure 3 & 4. For Viscosity $\eta = \eta_0(t/t_0 \times \rho/\rho_0)$ and Surface Tension $\gamma = \gamma_0(n_0/n \times \rho/\rho_0)$

RESULTS AND DISCUSSION

The Tomatoes fruits contained Lycopene extract in PtEt were quantities determined by using UV-Vis spectrophotometer at 503nm wavelength the CS1 has Lycopene 0.042783mg/100gm and another AS2 of 0.016227mg /100gm in which CS1 have more then quantities compared to AS2 now we determined the some physicochemical parameters such as viscosity and surface tension by BMS where viscosity of CS1 at 0.400467 N.sm⁻², and AS2 at 0.371466 N.sm⁻² and where Surface tension gives them their near-spherical shape, because a sphere has the smallest possible of CS1 at 20.59763 N.m⁻¹, AS2 at 22.87903 N.m⁻¹ the surface tension of samples CS1 < AS2 and pH values has compared between pure PtEt of 6.77 and both PtEt extracted Lycopene from Tomatoes was of CS1 at 6.25, AS2 at 6.60 on temperature at 20°C.





CONCLUSION

The Lycopene molecule has anticancer properties and much probability to reduce cancers. When we intake some varieties of tomatoes such as in food materials without cooking in which Lycopene is present but when we cook them, the heat temperature may degrade the Lycopene quantities [20]. According to these researches, the cherry type tomato has much quantities of Lycopene compared to other varieties of tomatoes.

ACKNOWLEDGMENT

The authors thank to Dean & Prof. Mansingh and Faculty, School of Chemical Sciences, CUG, Gandhinagar, for support.

REFERENCES

1. Dov Hartal Ph.D. and Lorri Damg” Industrial Production, Properties and Uses of Tomato Lycopene”, Additional Lyc-0-Mato* Information pp. 2-9, (2004).
2. M.J. Periago, F. Rincón, M.D. Agüera, G. Ros, Mixture approach for optimizing lycopene extraction from tomato and tomato products, *J. Agric. Food Chem.* 52, 5796–5802 (2004).
3. I. R. Bunghez, M. Raduly, S. Doncea, I. Aksahin And R. M. Ion “Lycopene Determination in Tomatoes by Different Spectral Techniques (UV-Vis, FTIR And HPLC)” *Journal of Nanomaterial’s and Biostructures* Vol. 6, p. 1349-1356(2011).
4. Stahl, W. and Sies, H., Lycopene: a biologically important carotenoid for humans? *Arch. Biochem. Biophys.* 336, 1–9(1996).
5. Schierle, J., Bretzel, W., Bühler, I., Faccin, N., Hess, D., Steiner, K., and Schüep, W., Content and isomeric ratio of lycopene in food and human blood plasma. *Food Chem.* 59, 459- 465.(1997).

6. Di Mascio, P., Aaiser, S., and Sies, H., Lycopene as the most elective biological carotenoid singlet oxygen quencher. *Arch. Biochem. Biophysics.* 274, 532-538(1989).
7. Gerster, H. The potential role of lycopene for human health. *J. Am. Coll. Nutr.* 16(2), 109–126(1997).
8. Giovannucci, E. “Tomatoes, tomato-based products, lycopene, and cancer: review of the epidemiological literature”. *J. Natl. Cancer Inst.* 91, 317–331(1999).
9. Giovannucci, E., Ascherio, A., Rimm, E. B., Stampfer, M. J., Colditz, G. S., and Willett, W. C., Intake of carotenoids and retinol in relation to risk of prostate cancer. *J. Natl. Cancer Inst.* 87, 1767–1776(1995).
10. Rao, A. V. and Agarwal, S., Bioavailability and in vitro antioxidant properties of lycopene from tomato products and their possible role in the prevention of cancer. *Nutr. Cancer* 31, 199–203(1998).
11. Sadler, G., Davis, J., and Dezman, D., Rapid extraction of lycopene and b-carotene from reconstituted tomato paste and pink grapefruit homogenates. *J. Food Sci.* 55, 1460–1461(1990).
12. Kimura, M. and Rodriguez-Amaya, D. B., Sources of errors in the quantitative analysis of food carotenoids by HPLC. *Arch. Latinoam. Nutr.* 49, 58S–66S (1999).
13. Halim Y, Schwartz SJ, Francis D, Baldauf NA, Rodriguaz-Saona LE, Direct determination of lycopene content in tomatoes (*Lycopersicon esculentum*) by attenuated total reflectance infrared spectroscopy and multivariate analysis. 89(5):1257-62(2006).
14. John W. Lehman, *Operational Organic Chemistry*; 4th Ed.
15. Sadasivan, S. and Manikhyam, A., *Biochemical methods for agricultural sciences*, Wiley Eastern Limited, New Delhi, pp.106-108 (1991).
16. Adsule, P. G. and Dan, A., Simplified extraction procedure in the rapid spectrophotometric method for lycopene estimation in tomato. *J. Food Sci. Technol.* 16, 216 (1979).
17. Wayne W. Fish, Penelope Perkins-Veazie, and Julie K. Collins A Quantitative Assay for Lycopene That Utilizes Reduced Volumes of Organic Solvents *Journal of Food Composition And Analysis* 15, 309–317(2002).
18. Zechmeister, L., A.L. Lerosen, W.A. Schroeder, A. Polgar and L. Pauling, Spectral characteristics and configuration of some stereo isomeric carotenoids including prolycopene and pro-carotene. *J. Am. Chem. Soc.*, 65: 1940-1951(1943)
19. Mansingh “Survismeter, 3-in-1 Instrument for Simultaneous Measurements of Surface Tension, Inter Facial Tension (IFT) and Viscosity” *Pak. J. Anal. Environ. Chem.* Vol. 8, No. 1& 2 (2007) 82 . 85

20. Anil Kumar et al (2014) "Effect of Temperature on Quantities of Lycopene in Some Replicate of Tomatoes" International Journal of Pharmacology & Toxicology / 4(3), 2014, 142-145.