

CHARACTERIZING THE ANTIOXIDANT AND ANTICANCER PROPERTIES OF SECONDARY METABOLITES FROM RED AND WHITE CABBAGES *BRASSICA OLERACEA L. VAR. CAPITATA*

Alaa A. Gaafar^{1*}, Hanan F. Aly², Zeinab A. Salama¹ and Khaled M. Mahmoud³

¹Plant Biochemistry Department, National Research Centre, El Behouth St., P.O. Box 12311, Dokki, Cairo, Egypt.

²Therapeutic Chemistry Department, National Research Centre, El Behouth St., P.O. Box 12311, Dokki, Cairo, Egypt.

³Pharmacognosy Department, National Research Centre, El Behouth St., P.O. Box 12311, Dokki, Cairo, Egypt.

Article Received on
15 April 2014,

Revised on 10 May 2014,
Accepted on 03 June 2014

*Author for Correspondence

Dr. Alaa Gaafar

Plant Biochemistry
Department, National Research
Centre, El Behouth St., P.O.
Box 12311, Dokki, Cairo,
Egypt

ABSTRACT

Vegetables belongs to *Brassicaceae* are rich in polyphenols, flavonoids, glucosinolates, vitamin C and their hydrolysis products which may have antioxidant and anticancer properties. This study was carried out to evaluate and differentiate the total phenolic content, total flavonoids, vitamin C and their antioxidant potential activities and cytotoxic effects between red and white cabbages after extraction with two different solvents (80% methanol and aqueous extracts). The total phenolic and total flavonoid contents varied from 29.13 to 11.44 mg/gdw and from 11.44 to 4.37 mg/gdw for red and white cabbage respectively that extracted with 80% methanol. Levels of total phenolic and flavonoids extracted with water were ranged from 21.38 to 10.36

mg/gdw and from 12.33 to 3.69 mg/gdw for red and white cabbages respectively. While, vitamin C ranged from 64.87 to 44.25 mg/100gfw. Red cabbage was the richest sources of dietary antioxidants, while their content was the lowest in white cabbage. Phenolic compounds present in the extracts showed antioxidant activities investigated using DPPH radical scavenging activity, iron chelating and reducing power. The extract of red cabbage exhibited a stronger ability to scavenge DPPH radical as compared to white cabbage. Iron chelating and reducing power activities were also stronger in red cabbage. Results of the HPLC analysis indicated that, among phenolic compounds, luteolin and rutin were the main constituents in red cabbage extracted with 80% methanol. Red and white cabbages were tested for their anti-cancer activity using three cancer cell lines; hepatocellular cell line

(HePG2), colon cell line (HCT116), breast cell line (HCF7) and a normal cell line (human dermal fibroblast cell line). Each extract is evaluated cytotoxicity inhibition effects. Red cabbage aqueous extract showed cytotoxic inhibition effects of 46.40, 31.20 and 22.40 % for HePG2, HCT116 and MCF-7, respectively. Red cabbage seems to be a very good source of dietary antioxidants possessing high antioxidant and anticancer activities. Thus, it could be concluded that, Brassica vegetables contain bioactive substances such as phenolic and vitamin C with potential cancer preventive effects.

Key words: Red cabbage, White cabbage, Antioxidant, Anticancer.

INTRODUCTION

Brassicaceae vegetables are an abundant source of health-promoting substances, which reduce the risk of diseases. Apart from anticancerogenic glucosinolates, they possess antioxidants of both hydrophilic (vitamin C, polyphenols) and hydrophobic phases (carotenoids, vitamin E), which can neutralize active oxygen species and quench free radicals. Some studies have been conducted to quantify the phenolic compounds, carotenoids, vitamin C, and antioxidant potential [1], [2]. It was found that, the majority of the antioxidant activity of Brassicaceae vegetables may be related to phenolic compounds such as flavonoids, isoflavone, flavones, anthocyanin, catechin and isocatechin [3].

The antioxidant properties were tested in many studies by using different approaches [4], [5]. The content of antioxidants depends on a lot of factors, especially on cultivars, stage of maturity and growing conditions [6]. The functional quality and antioxidant constituents of cabbage heads are strongly influenced by environmental factors and genetics. The ferric reducing antioxidant power (FRAP), Trolox equivalent antioxidant capacity (TEAC) and free radical scavenging activity (DPPH) assays are the three most frequently used for assessing the antioxidant activities [7]. Different botanical compounds gained attention as therapeutic agents that relieve pain and inflammation [8]. The objective of this study was focused on characterizing the antioxidant and anticancer properties of secondary metabolites from red and white cabbages.

MATERIALS AND METHODS

Chemicals and cells

Ascorbic acid, Folin–Ciocalteu reagents, Gallic acid, Quercetin, DPPH: 2,2-diphenyl-1-picrylhydrazyl, Ferrozine: (3-(2-pyridyl) -5, 6-bis- (4-phenylsulfonic acid)-1, 2, 4-triazine, BHT: Butyl Hydroxy toluene and, were from (Sigma Chemical Co., St. Louis, MO, USA).

Preparation of plant extract

The leaves were extracted. Briefly, 10 g of the dried powder from red and white cabbages were soaked with 100 ml of 80% methanol and water and shaking at room temperature for 48 h. The extracts were filtered and the extraction was repeated twice. The resulting methanolic and aqueous extracts were used for the determination of total phenolic, flavonoid, antioxidant activity and cytotoxic effects on different human cancer cell line (HePG2, HCT116 and MCF-7).

Total phenolic content (TPC)

The total phenolic content (TPC) of methanolic and water extracts of red and white cabbages were spectrophotometrically determined by FolinCiocalteu reagent assay using gallic acid for the preparation of calibration curve (20 – 120 mg/l) according to the method of [9]. A suitable aliquot (1 ml) of each extract or standard solution was added to 25 ml volumetric flask, containing 9 ml of distilled water. One milliliter of FolinCiocalteu's phenol reagent was added to the mixture and shaken. After 5 min. 10 ml of 7 % Na₂CO₃ solution were added to the mixture. The solution was diluted to 25 ml with distilled water and mixed. After incubation for 90 min. at room temperature, the absorbance was determined at 750 nm with spectrophotometer (Unicum UV 300) against prepared reagent as blank. A total phenolic content in samples was expressed as mg gallic acid equivalents (GAE)/g dry weight. All samples were analyzed in triplicates.

Total flavonoid content (TFC)

Total flavonoid content (TFC) of methanolic and water extracts of red and white cabbages were spectrophotometrically determined by the aluminum chloride method using quercetin as a standard [10]. One ml of extract or standard solution (quercetin, 20–120 mg/l) was added to 10 ml volumetric flask, containing 4 ml of distilled water. To the flask 0.3 ml 5 % NaNO₂ was added and after 5 min 0.3 ml 10 % Al Cl₃ was added. At 6th min, 2 ml 1M NaOH were added and the total volume was made up to 10 ml with distilled water. The solutions were mixed well and the absorbance was measured against prepared reagent blank at 510 nm by using spectrophotometer (Unicum UV 300). Total flavonoids in sample were expressed as mg quercetin equivalents (QE)/ g fresh weight. Samples were analyzed in triplicates.

Determination of vitamin C

Ascorbic acid content in fresh samples of red and white cabbage was measured colorimetrically according to [11].

Determination of free radical scavenging activity

Determination of free radical scavenging activity was performed spectrophotometrically as described by [12]. An aliquot of 0.5 ml of 0.1 mM DPPH[•] radical were added to a test tube with 1 ml of methanolic and aqueous extracts at different concentrations 5, 10, 20, 40 µg/ml. Methanol was used as blank. The reaction mixture was shaken vigorously at room temperature. , buthylatedhydroxyl toluene (BHT) was used as a positive control, and negative control contained the entire reaction reagent except the extracts. Then the absorbance was measured at 515 nm with spectrophotometer against blank (methanol pure). Lower absorbance of the reaction mixture indicated higher free radical scavenging activity. The capacity to scavenge the DPPH[•] radical was calculated using the following equation:

$$\text{DPPH}^{\bullet}\text{scavenging effect (Inhibition \%)} = [(A_c - A_s / A_c) \times 100].$$

Where A_c was the absorbance of the control reaction and A_s as the absorbance in the presence of Red and white extracts.

Determination of chelating effects on ferrous ions

Metal chelating effects on ferrous ions was carried out as described by [13]. One ml of methanolic and aqueous extracts, and or EDTA solution as a positive control at different concentrations (5, 10, 20, 40 µg/ml) were mixed with 0.1 ml of 2 mM $\text{FeCl}_2 \cdot 4\text{H}_2\text{O}$ and 0.2 ml of 5 mM ferrozine solution and 3.7 ml methanol or water were mixed in a test tube and reacted for 10 min, at room temperature. The absorbance was then measured at 562 nm. Mixture without extract was used as the control. A lower absorbance indicates a higher ferrous ion chelating capacity. The percentage of ferrous ion chelating ability was calculated using the following equation: Iron chelating activity (Inhibition %) = $[(A_c - A_s / A_c) \times 100]$

Where A_c was the absorbance of the control reaction and A_s as the absorbance in the presence of Red and white cabbage extracts.

Reducing power

The reducing power was assayed as described in [14]. Different concentrations (5, 10, 20, 40 µg/ml) of methanolic and aqueous extracts of red and white cabbages (1.0 ml) were mixed with 2.5 ml of phosphate buffer (50 mM, pH 7.0) and 2.5 ml of 1% potassium ferricyanide. The mixture was then incubated at 50°C for 20 min. After, 2.5 ml of trichloroacetic acid (10 %) were added to the mixture, centrifuged at 3000 rpm for 10 min. Finally, 1.25 ml from the supernatant was mixed with 1.25 ml of distilled water and 0.25 ml FeCl_3 solution (0.1%,

w/v). The absorbance was measured at 700 nm. The assays were carried out in triplicate and the results were expressed as mean values \pm standard deviations. Increased absorbance values indicate a higher reducing power. The extract concentration providing 0.5 of absorbance (EC_{50}) was calculated from the graph of absorbance at 700 nm against the extract concentration. BHT was used as standard.

Cytotoxic effect on human cell line (HePG2 – MCF7 – HCT116)

Cell viability was assessed by the mitochondrial dependent reduction of yellow MTT (3-(4, 5-dimethylthiazol -2-yl)-2, 5-diphenyltetrazolium bromide) to purple formazan [15].

Procedure: All the following procedures were done in a sterile area using a Laminar flow cabinet bio safety class II level (Baker, SG403INT, Sanford, ME, USA). Cells were suspended in RPMI 1640 medium for HePG2- MCF7 and HCT116. The media are supplemented with 1% antibiotic-antimycotic mixture (10,000U/ml Potassium Penicillin, 10,000 μ g/ml Streptomycin Sulfate and 25 μ g/ml Amphotericin B), 1% L-glutamine and 10% fetal bovine serum and kept at 37 °C under 5% CO₂. Cells were batch cultured for 10 days, then seeded at concentration of 10x10³ cells/well in fresh complete growth medium in 96-well microtiter plastic plates at 37 °C for 24 h under 5% CO₂ using a water jacketed Carbon dioxide incubator (Sheldon, TC2323, Cornelius, OR, USA). Media was aspirated, fresh medium (without serum) was added and cells were incubated either alone (negative control) or with different concentrations of sample to give a final concentration of (100 – 50 – 25– 12.5–6.25–3.125–0.78 and 1.56 μ g/ml). After 48 h of incubation, medium was aspirated, 40 μ l MTT salt (2.5 μ g/ml) were added to each well and incubated for further four hours at 37°C under 5% CO₂. To stop the reaction and dissolving the formed crystals, 200 μ l of 10% Sodium dodecyl sulphate (SDS) in deionised water was added to each well and incubated overnight at 37°C. A positive control which composed of 100 μ g/ml was used as a known cytotoxic natural agent who gives 100% lethality under the same conditions [16]. The absorbance was then measured using a micro plate multi-well reader (Bio-Rad Laboratories Inc., model 3350, Hercules, California, USA) at 595nm and a reference wavelength of 620nm. A statistical significance was tested between samples and negative control (cells with vehicle) using independent t-test by SPSS 11 program. DMSO is the vehicle used for dissolution of plant extracts and its final concentration on the cells was less than 0.2%. The percentage of change in viability was calculated according to the formula:

$$((\text{Reading of extract} / \text{Reading of negative control}) - 1) \times 100$$

A probit analysis was carried for IC₅₀ and IC₉₀ determination using SPSS 11 program.

Statistical Analysis

Data were statistically analyzed using Costat statistical package data according to [17].

RESULTS AND DESCUSSION

Total phenolic content, total flavonoids content and vitamin C

The quantitative contents of total phenolics, total flavonoids and vitamin C in the extracts of red and white cabbages are presented in (Table 1 and 2). The total phenolic and total flavonoid contents varied from 29.13 to 11.44 mg/gdw and from 11.44 to 4.37 mg/gdw for red and white cabbage respectively that extracted with methanol. Levels of total phenolic and flavonoids extracted with water were ranged from 21.38 to 10.36mg/gdw and from 12.33 to 3.69 mg/gdw for red and white cabbages respectively. While vitamin C ranged from 64.87 to 44.25 mg /100gfwof fresh red and white cabbage. Red cabbage methanolic extract was characterized by a higher content of total phenolics and total flavonoid.

Table 1. Total phenolic (TPC) and total flavonoids (TFC) in red and white cabbage

Extracts	Samples	TPC(mg*/g DW)	TFC(mg**/gDW)
Methanolic	Red cabbage	29.13 ^d ± 0.08	17.44 ^d ± 0.11
	White cabbage	11.36 ^b ± 0.02	4.37 ^b ± 0.14
Aqueous	Red cabbage	21.38 ^c ± 0.11	12.33 ^c ± 0.11
	White cabbage	10.36 ^a ± 0.04	3.69 ^a ± 0.09
LSD at 0.05		0.25	0.24

* As gallic acid. ** As quercetin.

All values are the means of three replicates and are significantly different at $p \geq 0.05$

Table 2. Vitamine C in red and white cabbages

Samples	Vit C(mg/100gF W)
Red cabbage	64.87 ^b ± 0.14
White cabbage	44.25 ^a ± 0.17
LSD at 0.05	0.14

The results are compatible with a study carried out by [18] who showed that the total phenolic content of the cabbage extract was 4.1 mg/g. Other authors reported that, in fresh white cabbage 0.4 mg total phenolics/g FM [19], 1.10-1.53mg total phenolics /g FM.[20], 0.36mg total phenolics/g FM[21].

Antioxidant activity

DPPH radical scavenging activity

All the cabbages extracts at the tested concentrations were capable of directly reaction with and quenching DPPH radicals Table (3). The red cabbage aqueous extract were found to have the highest DPPH scavenging activity ranged from (62.51 to 89.04 µg/ml), from (31.58 to 59.42 µg/ml) in white cabbage.

Table 3. DPPH Scavenging activity of red and white cabbage

Extracts	Samples	Inhibition %			
		5 µg/ml	10 µg/ml	20 µg/ml	40 µg/ml
Methanolic	Red cabbage	70.52 ^d ± 0.31	81.07 ^c ± 0.31	88.83 ^d ± 0.31	93.71 ^d ± 0.21
	White cabbage	35.50 ^b ± 0.33	42.75 ^a ± 0.26	57.80 ^b ± 0.36	70.24 ^b ± 0.26
Aqueous	Red cabbage	62.51 ^c ± 0.16	76.84 ^b ± 0.36	81.17 ^c ± 0.16	89.04 ^c ± 0.31
	White cabbage	31.58 ^a ± 0.31	42.82 ^a ± 0.31	51.03 ^a ± 0.41	59.42 ^a ± 0.31
BHT as Standard		71.55 ^c ± 0.27	81.99 ^d ± 0.79	89.86 ^e ± 0.36	94.40 ^e ± 0.36
LSD at 0.05		0.47	0.86	0.68	0.60

In comparison with white cabbage cultivar, antioxidant value was lower than that cabbage value used in other study [22]. The antioxidant activity of the plant can be attributed to flavonoids and polyphenolic compounds found in it [23]. Total phenol and flavonoid contents were significantly different between the studied cabbage cultivars ($p \leq 0.05$). However, the antioxidant capacity also depends on several other factors including genetics, environmental conditions, production techniques used, date of harvest and post-harvest storage conditions [24].

Fe-chelating activity of red and white cabbage

Red and white cabbages have significant ferrous ion chelating activity but in all cases it was significant lower than EDTA. The highest Fe chelating was found in red cabbage aqueous extract it ranged from 19.52 to 77.42 µg/ml and from 13.05 to 73.55 µg/ml in white cabbage (Table 4).

Table 4. Fe-chelating activity of red and white cabbage

Extracts	Samples	Inhibition %			
		5 µg/ml	10µg/ml	20µg/ml	40µg/ml
Methanolic	Red cabbage	35.90 ^c ± 0.49	54.70 ^d ± 0.55	82.95 ^d ± 0.28	90.03 ^d ± 0.28
	White cabbage	19.84 ^b ± 0.49	30.18 ^a ± 0.28	58.70 ^c ± 0.49	79.04 ^c ± 0.41
Aqueous	Red cabbage	19.52 ^b ± 0.41	38.64 ^c ± 0.47	56.91 ^b ± 0.21	77.42 ^b ± 0.47
	White cabbage	13.05 ^a ± 0.56	34.10 ^b ± 0.41	55.33 ^a ± 0.27	73.55 ^a ± 0.27
EDTA as Standard		49.48 ^d ± 0.41	68.65 ^e ± 0.55	83.67 ^e ± 0.49	93.43 ^e ± 0.28
LSD at 0.05		0.97	0.96	0.66	0.68

Antioxidant activity in *Brassica* species is correlated with vitamin C and total phenolic content [25]. In the previous investigations [26], ascorbic acid content in red and white cabbages was not differentiated as phenolic substances estimated in the present study, hence, high antiradical activity determined in red cabbage in comparison with the samples of white cabbage cultivars was probably due to phenolics, especially those of the phenylpropanoids and anthocyanins groups. According to [27], polyphenols possess the ideal chemical structure for free radical scavenging and seem to be more effective antioxidants than vitamins E and C. The phenolic acids, antiradical activity are positively correlated with the number of hydroxyl groups bonded to the aromatic ring [28]. In the present study, radical scavenging activity is measured by two methods: DPPH, Fe-chelating and power reducing reagents. In the case of DPPH, antiradical activity was higher as related to the Fe –chelating and power reducing method. A similar effect was described by [2] for Radical scavenging activity (RSA) estimated in white cabbage assays. The various results obtained by these three chemical methods might have been due to the reaction of different phenolics with free radicals of different chemical structures. Several studies have reported on the relationships between phenolic content and antioxidant activity [29],[30],[31]. Some authors found a correlation between the phenolic content and the antioxidant activity [32],[33],[28].

Reducing Power of red and white cabbage

Reducing power antioxidant activity was expressed as EC₅₀ (effective concentration at which the absorbance is 0.5 at 700 nm). Reducing power of methanolic and aqueous extracts of red and white cabbage in comparison with BHT as standard showed in (Fig 1 and Table 5).




Fig.1. Reducing power of 80% methanolic and aqueous extracts of red and whiteCabbage

Table 5.Reducing power activity of red and white cabbage

Extracts	Samples	EC ₅₀ µg/ml
Methanolic	Red cabbage	8.23 ^b ± 0.02
	White cabbage	9.87 ^e ± 0.13
Aqueous	Red cabbage	20.47 ^c ± 0.03
	White cabbage	28.49 ^d ± 0.14
BHT as Standard		4.09 ^a ± 0.10
LSD at 0.05		0.21

It was observed that methanolic extract of red and white cabbage had higher reducing power (EC₅₀ = 8.23 – 9.87 µg/ml) than aqueous extract (EC₅₀ = 20.47 – 28.49µg/ml), respectively. The reducing power activity may be due to the presence of reductones as they are electron donors and are capable of converting them into a more stable product and terminating the free radical reaction. The reducing power activity of the extracts examined in this study was similar with the results of [34] and [35] who reported that reducing power of broccoli extracts was used as a method for antioxidant activity investigation. The extract of white cabbage possessed also the stronger reducing power [36].

Identification of phenolics compounds by HPLC

The HPLC profile of crude methanolic extracts of Red and white cultivars extracted with methanol and water are illustrated in Table (6).

Table 6. Phenolic compounds of red and white cabbage cultivars (mg/100gD.W)

Phenolic compounds mg/ 100g DW	Methanolic		Aqueous	
	Red Cabbage	White Cabbage	Red Cabbage	White Cabbage
Pyrogallol	-	16.22	5.12	-
Gallic acid	-	3.07	3.67	3.10
Chlorogenic acid	1.59	0.43	2.29	0.79
Vanillic	-	0.38	0.32	0.77
Catechines	-	15.92	22.34	22.53
Caffeic acid	-	0.70	0.39	8.12
Rutin	58.36	5.76	13.48	4.49
<i>P</i> -Coumaric acid	-	-	10.72	1.47
Ferulic	7.82	1.45	1.30	5.19
Benzoic	14.40	1.80	2.15	6.34
Acacetin	8.06	3.32	1.78	4.55
Myricetine	7.61	-	-	0.82
Coumarin	5.36	-	-	0.79
Luteolin	119.65	10.82	38.56	2.95
Quercetin	36.33	7.98	-	-
Cinnamic acid	0.74	0.38	1.17	0.31
Genistein	2.57	-	2.22	0.27
Kaempferol	1.88	2.05	0.33	3.23

Methanolic extract enhanced phenolic compounds in both cultivars of cabbage as compared to water aquease extract. Methanolic extract of red cabbagecultivars showed higher content in most phenolic compounds especially Luteolin (119.65 mg/ 100gDW), Rutin (58.36 mg/100g DW), Qurectein (36.33 mg/100g DW),and Benzoic (14.40 mg/100g DW).

Whereas, Methanolic extract of white cabbage cultivar showed lower content in phenolic compounds especially in Pyrogallic acid (16.22mg/100gdw), Catchiness acid (15.92mg/100gDW), and Luteolin (10.82 mg/100gDW).

The least phenolic constituent was Acacetin, which was detected in red and white cabbage aqueous water extract.Caffeic acid and Kaempferol in red cabbage. Whereas, Ggenistein (0.27mg/100gDW) and Cinnamic 0.31mg/100 gDW).The results obtained showed significant quantitative variation in the polyphenolic profile as well as variation in the concentration of each individual compound. In this concern, [37] mentioned that phenolic compounds ranged between plants according to their genus species, varieties, cultivars, and types of fertilizer.

Cytotoxicity effect of red cabbage and white cabbage

Cancer is a global health problem with high morbidity and mortality and poses both economic and psychological challenges [38],[39].It is known that different cell lines might exhibit different sensitivities towards an antiproliferative compound, so the use of more than one cell line is therefore considered necessary in the detection of antiproliferative compounds.

Red and white cabbages were tested for their anti-cancer activity using three cancer cell lines hepatocellular cell line (HePG2), colon cell line (HCT116) and breast cell line (HCF7), beside normal cell line (human dermal fibroblast cell line). The cytotoxic activity data are presented in Table (7).

Table 7. Cytotoxicity effect of red and white cabbage

Extracts	Samples	Remarks % at 100 ppm		
		HePG 2	HCT116	HCF7
Methanolic	Red cabbage	35.8	26.7	19.8
	White cabbage	27.4	28.3	13.7
Aqueous	Red cabbage	46.4	31.2	22.4
	White cabbage	32.6	22.7	16.5

HePG2 :Human hepatocellular carcinoma cell line.

MCF7 : Human Caucasian breast adenocarcinoma.

HCT116:Human Colon Cancer Cells

Red cabbage aqueous extract exhibited a pronounced cytotoxic effect (46.40, 31.20 and 252.40 % for HePG2, HCT116 and MCF-7 respectively cell survival at 500 $\mu\text{g mL}^{-1}$)46.40, 31.20 and 252.40%, for HePG2, HCT116 and MCF-7 respectivelyas compared to that of white cabbage (32.605, 22.70, 16.50%).Cabbages were found to possess very potent inhibitory activities against all tested cell lines. These properties are in agreement with the known anticancer properties of cruciferous vegetables observed in both epidemiological and laboratory studies[40],[41],[42],[43].

Cruciferous vegetables contain several chemical compounds act as antioxidants or as inhibitors and/or inducers of phase I and phase II enzymes [44]. Phytonutrients in crucifers' vegetables work at a much deeper level and actually signal genes to increase production of enzymes involved in the detoxification.anti-carcinogenic actions of cruciferous vegetables are

attributed to their content of glucosinolate (GLS)[45]. Cruciferous vegetables contain sulforaphane, which has anticancer properties[46]. The anticancer effects of cruciferous vegetables may attribute to organic sulfur compounds (diallyl disulfide) and isothiocyanates, which had the ability to modulate expression/activity of antioxidative and phase 2 drug metabolizing enzymes and scavenging free radicals[41]. Also it was reported that administration of red cabbage extract to rats resulted in chemoprevention of liver and colon cancers induced by heterocyclic amine[43]. The same authors showed that breakdown products of glucosinolate such as isothiocyanates in red are responsible for the chemopreventive properties of cruciferous vegetables.

CONCLUSION

In conclusion, here in our study we demonstrated that the effects of white and red cabbages were found to have significant antioxidant and anticancer activity. These chemo preventive effects are likely related to the formation of organosulfur compounds following mechanical disruption of these vegetables, namely isothiocyanates from cruciferous. This may be a therapeutically insight in the treatment of cancers using *Brassicaceae* family.

REFERENCES

1. Nilsson J., Olsson K., Engqvist G., Ekvall J., Olsson M., Nyman M., Akesson B., (2006). Variation in the content of glucosinolates, hydroxycinnamic acids, carotenoids, total antioxidant capacity and low-molecular-weight carbohydrates in Brassica vegetables. *Journal of the Science of Food and Agriculture*, 86, 528-538.
2. Kusznierevicz, B, Bartoszek, A, Wolska, L, Drzewiecki, J, Gorinstein, S, Namiesnik, J, (2008). Partial characterization of white cabbages *Brassica oleracea var capitata f alba* from different regions by glucosinolates, bioactive compounds, total antioxidant activities and proteins. *LWT Food Science and Technology* 41, 1-9.
3. Wang, H., Cao, G.H., Prior, R.L., (1996). Total antioxidant capacity of fruits. *Journal of Agricultural and Food Chemistry*. 44, 701–705.
4. Liu, D., Shi, J., Colina Ibarra, A., Kakuda, Y. and Jun Xue, S. (2008). The scavenging capacity and synergistic effects of lycopene, vitamin E, vitamin C, and β -carotene mixtures on the DPPH free radical,” *LWT—Food Science and Technology*, 41, 1344–1349.

5. Zanfini, A., Corbini, G., La Rosa, C. and Dreassi, E. (2010). Antioxidant activity of tomato lipophilic extracts and interactions between carotenoids and α -tocopherol in synthetic mixtures, *LWT—Food Science and Technology*, 43, 67–72,
6. Hart, D.J, and Scott, K. J. (1995). Development and evaluation of an HPLC method for analysis of carotenoids in foods, and the measurement of the carotenoid content of vegetables and fruits commonly consumed in the UK. *Food Chemistry*. 54, 101–111.
7. Magalhaes LM, Segundo MA, Reis S, and Lima JL. (2008). Methodological aspects about in vitro evaluation of antioxidant properties. *Analytica ChimicaActa*, 613, 1-19.
8. Hernandez-Ortega M, Ortiz-Moreno A, Hernandez-Navarro MD, hamorro-Cevallos, G, Dorantes-Alvarez, L and Necoechea-Mondragón, H. (2012). Antioxidant, antinociceptive, and anti- Inflammatory effects of carotenoids extracted from dried pepper (*Capsicum annuum* L.). *Journal of Biomedicine and Biotechnology*, 2012, 1-10.
9. Singleton, V.L., and Rossi, J.A., (1965). Colorimetric of total phenolics with phosphomolibdic – phosphor tungstic acid reagents. *American Journal of Enology and Viticulture* 16, 144–158.
10. Zhishen, J., Mengcheng, T., and Jianming, W., (1999). The determination of flavonoid contents in mulberry and their scavenging effects on superoxide radicals. *Food Chemistry* 64, 555–559.
11. Bajaj, K.L. and Kaur, G. (1981). Spectrophotometric of L-ascorbic acid in vegetables and fruits. *Analyst*, 106, 117 – 120.
12. Chu, Y.F., Chang, C.L. and Hsu, H.F. (2000). Flavonoid content of several vegetables and their antioxidant activity. *Journal of Science of Food and Agriculture*. 80, 561–566.
13. Hsu, C., Chen, W., Weng, Y., Tseng, C., 2003. Chemical composition, physical properties, and antioxidant activities of yam flours as affected by different drying methods. *Food Chemistry*. 83, 85–92.
14. Kuda, T., Tsunekawa, M., Goto, H., & Araki, Y. (2005). Antioxidant properties of four edible algae harvested in the Noto Peninsula, Japan. *Journal of Food Composition and Analysis*, 18, 625–633.
15. Mosmann, T. (1983), Rapid colorimetric assays for cellular growth and survival: Application to proliferation and cytotoxicity assays. *Journal of Immunological Methods*. 65, 55–63.
16. Thabrew, M. I, Hughes, R. D. and McFarlane, I. G. (1997). Screening of hepatoprotective plant components using a HepG2 cell cytotoxicity assay. *Journal of Pharmacy and Pharmacology*. 49, 1132–1135.

17. Snedecor, G W., and Cochran, WG. (1989). Statistical methods.d.9th printing. Ames, IA: Iowa State Univ. Press.
18. Ismail, A., Marjam, Z. M., &Foong, C. W. (2004). Total antioxidant activity and phenolic content in selected vegetables. *Food Chemistry*, 87, 581–586.
19. Hassimotto, N.M., Genovese, M.I., and Lajolo ,F.M. (2005). Antioxidant activity of dietary fru, vegetables, and commercial frozen fruit pulps.*Journal of Agricultural and FoodChemistry*, 53, 2928–2935.
20. Kim, D.O.; Padilla-Zakour, O.I. and Grifths, P.D. (2004). Flavonoids and antioxidant capacity of various cabbage genotypes at juvenile stage.*Journal of Food Science*. 69, 685 – 689.
21. Marinova,D and. Ribarova, F (2005). Total phenolics and total flavonoids in Bulgarian fruits and getables, *Journal of the University of Chemical Technology and Metallurgy*, 40, 255-260.Chu, Y.F., Sun, J., Wu, X. & Liu, R.H. (2002).Antioxidant and antiproliferative activities of common vegetables.*Journal of Agricultural and Food Chemistry*. 50, 6910–6916.
22. Raghu KL, Ramesh CK, Srinivasa TR and Jamuna KS (2011). Total antioxidant capacity in aqueous extracts of some common vegetables. *Asian Journal of Experimental Biological Science*.2, 58-62.
23. Joseph B and Raj SJ (2010). Phytopharmacological properties of *Ficcusracemosa* Linn: An overview. *International Journal of Pharmaceutical Sciences Review and Research*.3, 134-8.
24. Dumas Y, Dadomo M, Di Lucca G, Grolier P. (2003). Effects of environmental factors and agricultural techniques on antioxidant content of tomatoes. *Journal of Science of Food and Agriculture*, 83, 369-82.
25. Hounsome N, Hounsome B, Tomos D, Edwards-Jones G. 2009. Changes in antioxidant compounds in white cabbage during winter storage.*Postharvest Biology and Technology* 52, 173-179.
26. Laszczak P., Szczeponek A., Wesołowska M., Leja M., (2007). Porównaniezawartościwybranychskładników w główkachkapustygłowiastejczzerwonej i białej.Nowaczyk P (ed.) Monography – Naturalna i indukowanazmienność w genetycznymdoskonaleniu roślinogrodniczych: 67-72.
27. Rice-Evans C.A., Miller N.J., and Paganga G., (1997). Antioxidant properties of phenolic compounds.*Trends in Plant Science*.2, 152-159.

28. Sroka Z., and Cisowski W., (2003). Hydrogen peroxidase scavenging, antioxidant and anti-radical activity of some phenolic acids. *Food and Chemical Toxicology*. 41, 753-758.
29. Podsędek a., 2007. Natural antioxidants and antioxidant capacity of Brassica vegetables: A review. *LWT* 40: 1-11.
30. Leja M., WyzGoLik G., mareczek a., 2005. Phenolic compounds of red cabbage as related to different forms of nutritive nitrogen. *Hort. Veg. Grow.* 24(3): 421-428.
31. Leja M., WyzGoLik G., kamińska i., 2007. Some parameters of antioxidant capacity of red cabbage as related to different forms of nutritive nitrogen. *Folia Hort.* 19(1): 15-23.
32. Kähkönen, M.P., Hopia, A.I., Vuorela, H.J., Rauha, J.P., Pihlaja, K., Kujala, T.S. and Heinonen, M., (1999). Antioxidant activity of plant extracts containing phenolic compounds. *Journal of Agricultural and Food Chemistry*, 47, 3954-3962.
33. Velioglu, Y. S., Mazza, G., Gao, L., and Oomah, B. D. (1998). Antioxidant activity and total phenolics in selected fruits, vegetables, and grain products. *Journal of Agricultural and Food Chemistry*, 46, 4113–4117.
34. Guo J.T., Lee H.L., Chiang S.H., Lin F.I., Chang C.Y., (2001). Antioxidant properties of the extracts from different parts of broccoli in Taiwan. *Journal of Food and Drug Analysis*, 9, 96–101.
35. Lin C.H., Chang C.Y., (2005). Textural change and antioxidant properties of broccoli under different cooking treatment. *Food Chemistry* , 90, 9–15.
36. Ciska, E., Karamac, M. and Kosińska, A. (2005). Antioxidant activity of extracts of white cabbage and sauerkraut. *Polish Journal of Food and Nutrition Science*. 14/55, 367–373.
37. Nakatani, N. 1997. Antioxidants from spices and herbs. In *Natural antioxidants: chemistry, health effects, and applications*, ed. F. Shahidi, 64–75. Champaign, IL: AOCS Press
38. Moyad, M.A. and Carroll, P.R. (2004). Life style recommendations to prevent prostate cancer, part II: time to redirect our attention? *Urologic clinics of North America* 31, 301–311.
39. Dossus, L., and Kaaks, R. (2008). Nutrition, metabolic factors and cancer risk”, *Best Practice & Research Clinical Endocrinology & Metabolism*. 22, 551–71.
40. Fleischauer, A. T., and Arab, L. (2001). Garlic and cancer: A critical review of the epidemiologic literature. *Journal of Nutrition*, 131, 1032S–1040S
41. Galeone, C., Pelucchi, C., Levi, F., Negri, E., Franceschi, S., Talamini, R., Giacosa, A., and La Vecchia C. (2006). Onion and garlic use and human cancer. *American Journal of Clinical Nutrition*, 84, 1027–1032.

42. Milner, J. A. (2001). Mechanisms by which garlic and allyl sulfur compounds suppress carcinogen bioactivation. Garlic and carcinogenesis. *Advances in Experimental Medicine and Biology*, 492, 69–81.
43. Talalay, P., and Fahey, J. W. (2001). Phytochemicals from cruciferous plants protect against cancer by modulating carcinogen metabolism. *Journal of Nutrition*, 131, 3027S–3033S.
44. Munday R, Munday CM. (2004). Induction of phase II enzymes by 3H-1, 2-dithiole-3-thione: doseresponse study in rats. *Carcinogenesis*. 25, 1721–1725
45. Brandi, G., Schiavano, G. F., Zaffaroni, N., De Marco, C., Paiardini, M., Cervasi, B. and Magnani, M. (2004). Mechanisms of action and antiproliferative properties of *Brassica oleracea* juice in human breast cancer cell lines. *Journal of Nutrition*, 135, 1503–1509.
46. Keum, Y.-S., Jeong, W.-S., & Kong, A. N. T. (2004). Chemoprevention by isothiocyanates and their underlying molecular signaling mechanisms. *Mutation Research*, 555, 191–202.