

## GENOTYPIC AND PHENOTYPIC CHARACTERIZATION OF SOME VIRULENCE GENES OF *KLEBSIELLA PNEUMONIAE* IN HILLA CITY

Massar A. M. AL-A`ammary\*, Jawad K. Al-Khafaji and Ifad K. Al-Shibly

Department of Microbiology, College of Medicine, University of Babylon, Iraq.

Article Received on  
12 June 2017,

Revised on 02 July 2017,  
Accepted on 23 July 2017

DOI: 10.20959/wjpr20178-8991

### \*Corresponding Author

Massar A. M. AL-  
A`ammary

Department of  
Microbiology, College of  
Medicine, University of  
Babylon, Iraq.

### ABSTRACT

*Klebsiella pneumoniae* is a gram-negative, non-motile, lactose fermenting, rod-shape organism. Although found in the microbiota, *K. pneumoniae* is frequently associated with hospital-acquired infections and can progress into severe illness. This study aims to detect some genes of virulence of *Klebsiella pneumoniae* isolated from different clinical samples in Hilla city, Iraq. A total (200) clinical samples were collected from different sources of patients from both sexes. Primary bacterial isolation was done, followed by phenotypic detection of some virulence factors, then molecular detection of virulence genes was done by genomic DNA extraction and Polymerase Chain Reaction (PCR). From a total of (200) clinical samples, only twenty isolates

(10%) of *Klebsiella pneumoniae* were identified according to the cultural characteristics and biochemical tests. Most isolates were in male gender, mean age of isolation was 14.9 years. Phenotypically all *Klebsiella pneumoniae* isolates were positive for urease, siderophore, colonization factors antigens and capsule production. Similarly, all isolates had strong ability for stone formation after incubation in normal sterile urine. While only 40% of isolates had high capacity for biofilm formation. Genotypic study revealed that among total of 20 local isolates of *Klebsiella pneumoniae*, all of the twenty isolates (100%) were negative for *mag A* gene, *Kfu* gene was detected in 5 isolates of *Klebsiella pneumoniae* (25%) and absent in 15 (75%) isolates. While *Uge* gene was found in two isolates (10%) only.

**KEYWORDS:** *Klebsiella pneumoniae*, *mag A* gene, *Kfu* gene, *Uge* gene.

## INTRODUCTION

*Klebsiella pneumoniae* is encountered as pathogen for humans and other mammals, colonizing different clinical sites including the gastrointestinal tract, skin, and nasopharynx; it is also found in various environmental niches (soil, water, etc.).<sup>[1]</sup> The pathogenicity of *K. pneumoniae* is multifactorial including LPS, capsule, urease, adhesins, outer-membrane proteins and biofilms.<sup>[2]</sup> Mucoviscosity-associated gene A (*magA*), has been identified in *Klebsiella pneumoniae*.<sup>[3]</sup> The *magA* is detected in a vast majority of *K. pneumoniae* liver abscess isolates and is associated with hypermucoviscosity (HV) and resistance to killing by human serum and phagocytosis.<sup>[4]</sup> The *kfu* gene encodes an iron-uptake system that is associated with a hypermucoviscosity phenotype and increased virulence.<sup>[5]</sup> While *K. pneumoniae* typically expresses both smooth lipopolysaccharide (O antigen) and capsule polysaccharide (K antigen) on its surface and both LPS and capsule contribute to the pathogenesis of this species. A single mutation in a gene that codes for a UDP-galacturonate 4-epimerase (*uge*) renders a strain with the O–K– phenotype (lack of capsule and LPS without O antigen molecules and outer core oligosaccharide).<sup>[6]</sup> This study aims to detect some genes of virulence of *Klebsiella pneumoniae* isolated from different clinical samples in Hilla city, Iraq.

## MATERIALS AND METHODS

A total (200) clinical samples were collected from different sources of patients from both sexes who attended or admitted to the main three hospitals in Hilla city /Babylon province (General Teaching hospital, Medical Teaching hospital of Merjan and Babylon hospital for Maternity and Children) for a period from September 2016 to December 2016. The different clinical specimens (sputum, sinus, tonsils, urine and vagina) were taken by a sterile containers, then stored in a cool place until transport to the laboratory. Primary bacterial isolation was done by streaking a loopful of culture on MacConkey agar for primary selection of pathogenic *Klebsiella pneumoniae*. The plates were incubated overnight at 37°C. The developed colonies which showed characteristic growth, colors, mucoid phenotype were transferred to a new plate for further tests.

Phenotypic detection of virulence factors was done individually for each virulence factor including capsule production by Indian ink test, Urease production by inoculating urea medium, Colonization factor antigen by the slide agglutination technique, Siderophore / Hemolysin production by inoculating a blood agar medium, Biofilm Production by tissue

culture plate method (TCP) assay and In Vitro Stone Formation by inoculating a fresh urine sample.


Molecular detection of virulence genes was done by Genomic DNA extraction followed by Polymerase Chain Reaction (PCR). The primers, which were used in PCR assay in this study with their sequences, and the DNA amplicon size the PCR conditions was carried out through the thermal cycling profile listed in table (1).

**Table 1: the primers used in the present study**

Markers	Oligonucleotide sequence 5`-3`	Size of amplicon	Thermal cycler conditions	Reference
<i>MagA</i> -F <i>MagA</i> -R	GGTGCTCTTTACATCATT GC GCAATGGCCATTTGCGT TAG	1280 pb	94C° for 1 min. 94C° for 30 sec. 59C° for 45 sec (30 cycle) 74C° for 2 min. 72C° for 6 min.	Fang <i>et al.</i> (2004)
<i>Kfu</i> -F <i>Kfu</i> - R	GAAGTGACGCTGTTTCT GGC TTTCGTGTGGCCAGTGA CTC	797 pb	94C° for 5min. 94C° for 60 sec. 54C° for 45 sec.(35 cycle) 72C° for 60 sec. 72C° for 7min.	Ma <i>et al.</i> , (2005)
<i>Uge</i> - F <i>Uge</i> - R	TCTTCACGCCTTCCTTCA CT GATCATCCGGTCTCCCTG TA	534 pb	94C° for 5 min. 94C° for 60 sec. 54C° for 45 sec.(35 cycle) 72C° for 60 sec. k72C° for 7 min.	Regue <i>et al.</i> , (2004)

## RESULTS

From a total of (200) clinical samples, only twenty isolates (10%) of *Klebsiella pneumoniae* were identified according to the cultural characteristics and biochemical tests as presented in Figure (1).




**Figure (1): Distribution of *Klebsiella pneumoniae* isolates out of total clinical samples (N=200).**

In the present study the *Klebsiella pneumoniae* isolates were distributed regarding different site of collection as the following: 8(40%) isolates from sputum, 5(25%) isolates from urine, 3(15%) isolates from tonsil, 2(10%) isolates from sinus, and also 2(10%) isolates from vagina as showed in Table (2).

**Table (2): Distribution of *Klebsiella pneumoniae* isolates according to the site of infection (N=20).**

Site of infection	No. of specimens	No. of <i>K.pneumoniae</i>	Percentage (%)
Sputum	70	8	40
Tonsil	55	3	15
Sinus	25	2	10
Vagina	20	2	10
Urine	30	5	25
<b>TOTAL</b>	<b>200</b>	<b>20</b>	<b>100</b>

In the present study the gender distribution of the *Klebsiella pneumoniae* isolation was studied and it was found that, *Klebsiella pneumoniae* in males formed 12 isolates(60%) and 8 isolates(40%) in the females as presented in Figure (2).



**Figure (2): Distribution of *Klebsiella pneumoniae* isolates out of total clinical samples according to gender of patients**

In the existent study the relationship between *Klebsiella pneumoniae* isolation rate and age of patients was investigated as presented in Table (3). The age-groups of patients who involved in present study was ranged from one year to thirty years old. Our results showed that the mean age of isolation was 14.9 years with standard deviation of 10.96.

**Table (3): Mean and standard deviation of the age of *Klebsiella pneumoniae* patients (N=20).**

Variable	Means $\pm$ SD	Range
Age (years)	14.9 $\pm$ 10.96	(1-30)

Concerning Urease production, The results in the present study showed that all *Klebsiella pneumoniae* isolates (100%) were positive for urease production.

In the present study, capsular phenotype was investigated among *Klebsiella pneumoniae* isolates, the results showed that all *Klebsiella pneumoniae* isolates (100%) were positive for capsule production.

The results of this study about siderophore production revealed that (100%) of *Klebsiella pneumoniae* isolates were able to produce siderophore (in presence of dipyrilidil) and none was able to produce hemolysin enzyme as shown in Table (4).

**Table (4): The siderophore and hemolysin production by *K.pneumoniae*.**

No. of <i>Klebsiella pneumoniae</i> isolates	Hemolysin production no.(%)	Siderophore production no.(%)
20	0(0%)	20(100%)

The colonization factors antigens study results through this work revealed that all isolates (100%) of *Klebsiella pneumoniae* that have been isolated from urine, sputum, vagina, sinus, and tonsil samples have the ability to express the colonization factor antigens (CFA/I, and CFA/III) as shown in Figure (3).




**Figure (3): Colonization Factor Antigens of *Klebsiella pneumoniae***

Biofilm formation ability of *Klebsiella pneumoniae* was investigated in this study and the results showed that 8 isolates (40%) had high capacity of biofilm formation due to strong adherence, 7 isolates (35%) had moderate capacity of biofilm formation, while only 5 isolates (25%) were weak producers of biofilm, the results presented in table (5).


**Table: (5) Biofilm formation of *Klebsiella pneumoniae*.**

Degree of Biofilm formation	No. of isolates	%
Strong	8	40
Moderate	7	35
Weak	5	25


In the present study, the stone formation ability has been investigated among urinary tract infection isolates of *Klebsiella pneumoniae* results showed that all isolates had strong ability for stone formation after incubation in normal sterile urine samples as shown in Figure (4).

**Figure (4): The stone forming ability of *K. pneumoniae* isolated from urine samples.**


Three selected genes were investigated molecularly in this study to explore their role in the pathogenesis of *Klebsiella pneumoniae* associated diseases. These genes were *mag A* gene (mucoviscosity associated gene A), *Kfu* gene (iron uptake system), and *Uge* gene (smooth LPS and O-antigen and K-antigen) (UDP-galacturonate 4-epimerase gene) by PCR technique. Among total of 20 local isolates of *Klebsiella pneumoniae*, the molecular detection for *mag A* gene by PCR revealed that all of the twenty isolates (100%) were negative for this gene. Regarding *Kfu* gene it was detected in 5 isolates of *Klebsiella pneumoniae* (25%) and absent in 15 (75%) isolates. While *Uge* gene results showed that only two isolates (10%) were positive for this gene out of the total twenty as presented in Figure (5), Figure (6), and Figure (7).

**Figure (5): Distribution of the virulence gene (*Mag A*, *Kfu* and *Uge*) among the *Klebsiella pneumoniae* isolates (N=20).**





**Figure (6):** Agarose gel electrophoresis of PCR product of (*Kfu*) gene amplicon product in *Klebsiella pneumoniae*. Lane 1-20 refer to isolate`s number.



**Figure (7)** Agarose gel electrophoresis of PCR product of *Uge* gene amplicon product in *Klebsiella pneumoniae* lane 1-12 refer to isolate`s number.

The distribution of virulence genes (*kfu*, *uge*) according to the site of collection among *Klebsiella pneumoniae* isolates had been studied, results presented in Table (6) and Table (7).

**Table (6):** Distribution of *Kfu* gene by source of infection (N=20).

Source of collection	<i>Kfu</i> gene		Total Number (%)
	Positive Number (%)	Negative Number (%)	
Sputum	3(15%)	5(25%)	8(40%)
Vagina	0(0.0%)	2(10%)	2(10%)
Sinus	1(5%)	1(5%)	2(10%)
Tonsil	1(5%)	2(10%)	3(15%)
Urine	0(0.0%)	5(25%)	5(25%)
TOTAL	5(25%)	15(75%)	20(100.0%)

**Table (7): Distribution of *Uge* gene by source of infection (N=20).**

Site of collection	<i>Uge</i> gene		Total Number (%)
	Positive Number (%)	Negative Number (%)	
Sputum	0(0.0%)	8(40%)	8(40%)
Vagina	0(0.0%)	2(10%)	2(10%)
Sinus	0(0.0%)	2(10%)	2(10%)
Tonsil	0(0.0%)	3(15%)	3(15%)
Urine	2(10.0%)	3(15%)	5(25%)
<b>TOTAL</b>	<b>2(10.0%)</b>	<b>18(90.0%)</b>	<b>20(100.0%)</b>

## DISCUSSION

This study results about the percentage of *Klebsiella pneumoniae* isolation rates were confirmed by the findings obtained by<sup>[7]</sup> and<sup>[8]</sup> who illustrated that immunocompromised patients were more frequently colonized by *Klebsiella* spp., and agreed with the findings obtained by<sup>[9]</sup> who instituted that *Klebsiella* spp. were responsible for many nosocomial infections in adults.

Results about the demographic data distribution were agreed with many previous studies such as that established by<sup>[10,11]</sup>, who found that *K. pneumoniae* was the commonest member of *Klebsiella* spp. that can cause severe infections. Among all patients diagnosed with *K. pneumoniae* pneumonia,<sup>[12]</sup> established that there was a male predominance thus more strengthening the results obtained in this current work.

A gender distribution difference was observed in this study, results came in agreement with<sup>[13]</sup> who reported that males are more susceptible to infection with *Klebsiella* spp. than females. Otherwise, these results did not agreed with other study which found that the percentage of *Klebsiella pneumoniae* in female was higher than male<sup>[14]</sup>

Results about the age distribution in the present study coming inside with research results of<sup>[15]</sup> who found a dominance of infection in age group range 16-30 years. Although other study found the percentage of *Klebsiella pneumoniae* is high in ages with 45-60 years.<sup>[16]</sup> Also, results about urease production agreed with<sup>[17]</sup>, who found that urease production is considered as an important virulence factor in bacterial pathogenicity because urease material elevated the pH due to the ammonia generated by this enzyme has important ramifications for medicine. Urease is a virulence factor in pathogenic bacteria that cause gastric ulceration, urinary stone formation, pyelonephritis, and other human health-related problems.<sup>[18]</sup>



The results about capsule production in the present study agreed with<sup>[17]</sup>, who found that capsule production is considered as an important virulence factors in bacterial pathogenicity because capsular material forms thick bundles of fibrillous structures covering the bacterial surface in massive layers.

Likewise, this study results about siderophore and heamolysin production are similar to those reported by<sup>[19]</sup> and<sup>[20]</sup>, who found that *Klebsiella* spp. have the ability to uptake iron through production siderophore but not hemolysin production.

All *Klebsiella pneumoniae* isolates (100%) produced CFA/III by hemagglutination of red blood cell group A in presence of tannic acid. This result agrees with finding obtained by<sup>[21]</sup> who finds that *Klebsiella* mediates mannose-resistant hemagglutination tannin treated erythrocyte.

The results of the present study about biofilm production came in agreement with<sup>[22]</sup> who found that a high value of biofilm formation is strong in urine sample, one of the most commonly studied properties of biofilms is their increased resistance to the effect of antibiotics.<sup>[23]</sup> And also this results agreed with<sup>[24]</sup> who found the biofilm formation should be considered in antimicrobial therapy.

Urinary stones may develop as a result of urinary tract infections. The latter plays an important role in urolithiasis. Whereas all non-infectious urinary stones originate because of metabolic disturbances or through changes so far unknown (for example in kidney tissue), coming inside with this study finding about the ability of *Klebsiella pneumoniae* for stone formation other study found that the commonest organisms isolated from urinary stone were *Klebsiella* in 12%.<sup>[25]</sup>

The results obtained in the current study about mag A gene among *Klebsiella pneumoniae* means that there are other means play a role in formation the mucoid viscosity. Although mag A gene is highly important for *Klebsiella pneumoniae* which confirm the bacteria mucoid viscosity, but it is prevalence and among *Klebsiella pneumoniae* local isolates is not high.<sup>[26]</sup> Many studies have indicated that mag A is not prevalent among *K. pneumoniae* isolates.<sup>[27],[18]</sup>

The *Kfu* gene which codes for an iron uptake system is a putative pathogenic gene, significantly associated with the virulent hypermucoviscosity phenotype and purulent tissue

infections.<sup>[28]</sup> Results about the prevalence of *Kfu* gene among *Klebsiella pneumoniae* in the existing study agreed with<sup>[29]</sup> who found the percentage of *kfu* gene is 25%. And these results disagreement with<sup>[30]</sup> the percentage is slightly higher than our study 66%.

Among the total of 20 isolates of *K. pneumoniae*, the molecular technique for *Uge* gene as done by PCR and detected in only two isolates (10%). Although this results disagree with<sup>[31]</sup> who found the percentage of *uge* gene higher than our study.

## CONCLUSION AND RECOMMENDATION

The conclusions that can be extrapolated by this study are that, among this study the local isolates of *Klebsiella pneumoniae* are a common among males than females especially in age group 14.9. Stone formation capacity of *Klebsiella pneumoniae* isolates may acts as contributing factor in their virulence. The *mag A* gene is absent among *Klebsiella pneumoniae*, other alternative virulence factor may play more important role in our local isolates. While the prevalence of *Kfu* gene and *uge* gene is relatively low among local *Klebsiella pneumoniae* isolates.

## REFERENCES

1. Lederman E.R, Crum N.F. Pyogenic liver abscess with a focus on *Klebsiella pneumoniae* as a primary pathogen: an emerging disease with unique clinical characteristics. *Am. J. Gastroenterol*, 2005; 100: 322–331.
2. Min W.Y, and Xuefing L.I. *Klebsiella pneumoniae* and *Pseudomonas aeruginosa*. *Molecular Medical Microbiology*, 2015; 1547-1564.
3. Fang CT, Chen YC, Chang SC, Sau WY, Luh KT. *Klebsiella pneumoniae* meningitis: timing of antimicrobial therapy and prognosis. *QJM*, 2000; 93: 45–53.
4. Lee H.C, Chuang YC, Yu WL, Lee NY, Chang CM, Ko NY. Clinical implications of hypermucoviscosity phenotype in *Klebsiella pneumoniae* isolates: association with invasive syndrome in patients with community-acquired bacteraemia. *J. Int. Med.*, 2006; 259: 606–614.
5. Hsu CR, Lin TL, Chen YC, Chon HC, Wang JT, The role of *klebsiella pneumoniae* *rpm A* in a capsular polysaccharides synthesis and virulence revisited. *Microbiology*, 2011; 157: 3446-3457.
6. Migueleagué, Beatriz Hita<sup>1</sup>, Nuria Piqué<sup>1</sup>, Luis Izquierdo, Susana Merino, Sandra Fresno, Vicente Javier Benedí<sup>1</sup>, and Juan M. Tomás. A Gene, *uge*, Is Essential for *Klebsiella pneumoniae* Virulence. *Infect Immun*, 2004; 72(1): 54-61.

7. Namratha, K.G., *et al.* Characterization and antibiogram of *Klebsiella* spp. isolated from clinical specimen in a rural teaching hospital. *Sch. J. App. Med. Sci.*, 2015; 3(2): 878–883.
8. Abhilash, K.P.P., Veeraraghavan, B., Abraham, O.C. Epidemiology and outcome of bacteremia caused by extended spectrum beta-lactamase (ESBL)-producing *Escherichia coli* and *Klebsiella* spp. in a tertiary care teaching hospital in South India. *J. Assoc. Physicians India*, 2010; 58: 13–17.
9. Tu, YC, Lu, MC, Chiang MK, Huang SP, Peng HL, Chang HY, *et al.* Genetic requirements for *Klebsiella pneumoniae*-induced liver abscess in an oral infection model. *Infect Immun*, 2009; 11: 21-9.
10. Ravichitra<sup>1</sup>,K.N. P.Hema Prakash, S. Subbarayudu and U.Sreenivasa Rao., Isolation and antibiotic sensitivity of *Klebsiella pneumoniae* from pus, sputum and urine samples, 2014; 3(3): 115-119.
11. Biradar, S. and C. Roopa. Isolation and Antibiogram of *Klebsiella* species from Various Clinical Specimens, 2015; 4(9): 991-995.
12. Lin,Y; Yu-Ping Wang, Fu-Der Wang and Chang-Phone Fung. Community-onset *Klebsiella pneumoniae* pneumonia in Taiwan: clinical features of the disease and associated microbiological characteristics of isolates from pneumonia and nasopharynx. *Front. Microbiol*, 2015; 18(8): 122-129.
13. Janda. J.M, & Abbott, S.I. The genera *klebsiella* and *Roultella*. *The Enterobacteraceae* (2<sup>nd</sup> edition., 2006; 115-129). Washingtonia, USA: ASMpress).
14. Shashidhar, T.P.K. Murthy, K.G. Girish, B. Manohar. Grinding of coriander seeds: modeling of particle size distribution and energy studies *Part. Sci. Technol.*, 2013; 31: 449-457.
15. Osazuwa F, 1Mordi RM, Osazuwa E, Taiwo SS, Alli OAT, Ogbolu DO, Akanni EO, Anukam; *Klebsiella* has taken lead among uropathogens in University of Benin Teaching Hospital, Benin City, Nigeria-An observation. *New York Science Journal*, 2010; 3(11): 61-64.
16. Russo, T. A. & Johnson, J. R. Medical and economic impact of extraintestinal infections due to *Escherichia coli*: focus on an increasingly important endemic problem. *Microbes Infect*, 2003; 5: 449–456.
17. Power, M.L; J. Littefield – Wyer, D.M. Gorden, D.A. veal and M.B. Slade. Phenotypic and genotypic characterization of Encapsulated *Escherichia coli* isolated from blooms in two Australian lakes. *Env. Microbiol*, 2004; 7(5): 631-640.

18. Rivero, A; Eric, G; David, A; David, B.H; Tom, C. K2 serotype *Klebsiella pneumoniae* causing a liver abscess associated with infective Endocarditis. *J. Clinic. Microbiol*, 48: 639-641.
19. Trakkanen, A.M; Allen, B.L, Williams, P.H; Kauppi, M; Haahtela, K; Siitonen, A. Fimbriation, capsulation and iron scavenging system of *Klebsiella* strains associated with human urinary tract infection. *Infect. Immune*, 1992; 60(3): 1187-1192.
20. Bingen, E; Picard, B; Brahimi, N; Mathy, S; Desjardins, P; Elion, J. and Denamur, E. Phylogenetic analysis of *Escherichia coli* strains causing meningitis suggests horizontal gene transfer from a predominant pool of highly virulent B2 group strains. *J. infect. Dis.*, 1998; 177: 642-650.
21. Trakkanen, A.M; Allen, B; Westerlund, B; Holthofer, H. Kuusela, P; Risteli, L; Clegg, S. and Korhonen, T.K. Type V collagen as target for type-3 fimbriae, Enterobacterial adherence. *Mol. Microbial*, 1990; 4: 1353-1361).
22. Kimia Seifi Hossein Kazemian, 2 Hamid Heidari, 3 Fereshteh Rezagholizadeh, 2 Yasaman Saeed, 4 Fariba Shirvani, 1 and Hamidreza Hour. Evaluation of Biofilm Formation Among *Klebsiella pneumoniae* Isolates and Molecular Characterization by ERIC-PCR. *Jundishapur J Microbiol*, 2016; 9(1).
23. Liaqat I, Sumbal F, Sabri AN. Tetracycline and chloramphenicol efficiency against selected biofilm forming bacteria. *Curr. Microbiol*, 2009; 59(2): 212-220.
24. Samia Bellifa, Hafida Hassaine, Damien Balestrino, Nicolas Charbonnel, Imane M'hamedi, Ibtissem Kara Terki, Merieme Lachachi, Wafae Didi and Christiane Forestier. Evaluation of biofilm formation of *Klebsiella pneumoniae* isolated from medical devices at the University Hospital of Tlemcen, Algeria. *African Journal of Microbiology Research*, 2013; 7(49): 5558-5564.
25. Alkadasi, b, Gamal A. A-Amerib, Samira Hanashb, Wahid A. M. Alib, Abdulrahman S. Najic, Nada A. Saifb, Abdulsalam Alshami and Abdubaset A. Zaidb. Incidence of renal stone disease among urinary tract infection patients and antimicrobials susceptibility. *Advances in Applied Science Research*, 2014, 5(3): 309-314
26. Abdul Razzaq, M. S; Al Khafajii, J.K; Al-Maamory E. H. K. Genotyping and Detection of Some Virulence Genes of *Klebsiella pneumoniae* Isolated from Clinical Cases. *Medical. J. of Babylon*, 2013; 10(2): 394-395.
27. Regue, M., Hita, B., Pique, N., Izquierdo, L., Merino, S., Fresno, S., Ben-edi, V. J., and Tomas J. M.: A Gene, *uge*, Is Essential for *K. pneumoniae* Virulence. *Infect. Immun*, 2004; 54-61.

28. Ma, L. C., Fang, C. T., Lee, C. Z., Shun, C. T. and Wang, J. T. Genomic heterogeneity in *K. pneumoniae* strains is associated with primary pyogenic liver abscess and metastatic infection. *J. Infect. Dis.*, 2005; 192: 117–128.
29. Aher T, Roy A, Kumar P. Molecular Detection of Virulence Genes Associated with Pathogenicity of *Klebsiella* spp. isolated from the respiratory tract of apparently healthy as well as sick goats. *Isr. J.Vet. Med.*, 2012; 67(4): 249-52.
30. Weldhagen GF, Poirel L, Nordmann P. Ambler class A extended-spectrum  $\beta$ -lactamases in *Pseudomonas aeruginosa*. *Antimicrob Agents Chemother*, 2003; 47: 2385–2392.
31. Aljanaby, A.A.J and Alhasani, A.H.A. Virulence factors and antibiotic susceptibility patterns of multidrug resistance *Klebsiella pneumoniae* isolated from different clinical infection. *African Journal of Microbiol. Res.*, 2016; 22(10): 829-834.