

EVALUATION OF ANTIPILEPTIC ACTIVITY OF METHANOLIC EXTRACT OF OPUNTIA FICUS INDICA FLOWER IN MICEKalbant Kumar*¹ and N.C. Nagalakshmi¹¹Mallige College of Pharmacy # 71, Silvepura, Chikkabanavara Post Bangalore: 560090.Article Received on
31 May 2017,Revised on 21 June 2017,
Accepted on 11 July 2017

DOI: 10.20959/wjpr20178-7853

Corresponding Author*Kalbant Kumar**

Mallige College of

Pharmacy # 71, Silvepura,

Chikkabanavara

Post Bangalore: 560090.

ABSTRACT

Opuntia ficus indica flowers (L) commonly known as cactus pear Flower, Indian fig Flower, belongs to the Cactaceae family. In the present study, Methanolic extract of *Opuntia ficus indica* flowers (MEOF) was tested for anticonvulsant activity in albino mice, using different convulsive models. The Methanolic extract of *Opuntia ficus indica* flowers produced significant anticonvulsant activity in all models studied. The present study also revealed that MEOF increase the concentration of nor-adrenaline and dopamine in brain of mice which attributed to the significant protection against MES induced seizures. Significant results were observed in the estimated parameters thereby justifying the use of this medicinal plant in the treatment of

epilepsy.

KEYWORDS: Epilepsy, *Opuntia ficus indica*, maximal electroshock, pentylenetetrazole, strychnine, seizures, Nor-adrenaline, Dopamine.

INTRODUCTION

Epilepsy is one of the central nervous system disorder that affect wide range of people throughout the world.^[1] The exact cause of epilepsy is not known however, it is believed that it occurs due to uncontrolled high frequency discharge of impulses by a group of neurons in the brain.^[2] Epilepsy is defined as recurrent and unprovoked epileptic seizures.^[3] The loss of consciousness, violent spasmodic contractions of skeletal muscles (convulsions) and autonomic hyper activity is usually associated with epilepsy.^[4]

The various mechanisms which involves the development of seizures either normal or pathologic brain are, 1) Diminution of inhibitory mechanism (especially synaptic inhibition

due to GABA) 2) Enhancement of the excitatory synaptic mechanism (especially those mediated by NMDA). 3) Enhancement of endogenous neuronal burst firing (usually by enhancing voltage dependent calcium currents).^[5]

There are many mechanism of action involved overcoming to the seizure like enhancement of GABA-mediated inhibition, suppression of rapid repetitive firing, reduction of current through T-type Ca^{++} channels, reduction of excitatory glutaminergic neurotransmission.^[6]

In this modern era of medicine there are 20 medications suggested by the Food and Drug Administration for the use of treatment of epileptic seizures. But, these drugs possesses various limitations in the form of serious side effects.^[7] Thus due to prevalent side effects and tolerance development of the presently available treatment for epilepsy, it is a necessary to develop a new drug without or less side effect.^[8,9]

MATERIALS AND METHOD

Plant material

The flowers of *Opuntia ficus indica* were collected in spring (Jun to September) from Puranpur, Pilibhit district, Bareilly, Uttar Pradesh, India. The plant was identified and authenticated by Dr. N. M. Ganesh Babu, senior botanist at FRLHT (Foundation for Revitalisation of Local Health Traditions) Jarakabande Kaval; post Attur, Yehalanka and Bangaluru (560106). A herbarium specimen was preserved in the college museum for future reference.

Extraction procedure^[10-12]

The flowers of *Opuntia ficus indica* was dried under shade and then powdered with a mechanical grinder. The powder was stored in an air tight container for further use. The coarse powder was extracted directly with 65% methanol (hydro-Methanolic extract) after defating with petroleum ether by Soxhlet extraction for 72 hours. The extracts were concentrated under reduced pressure and stored in a dissector until further use.

Animals

Swiss albino mice (20-25g) were selected and procured from Bioneds, Daverhosaholli, Sonapura hobli, Nelamangala (Tq), Tumkur (Dist.), Karnataka, India. They were acclimatized for one week under standard laboratory conditions (20-25°C and elimination cycle set to 12 hour light and 12 hour in dark) and fed with standard Amrut brand pellet diet and UV purified and filtered water was provided *ad libitum* in polypropylene bottles with

stainless steel sipper tubes. Each group contains six animals and was used for the mentioned research activities. The study protocol was approved by the Institutional Animal Ethics Committee (IAEC), Office of institutional animal ethical committee (IAEC) of Mallige College of pharmacy, Bangalore (IAEC 1432/PO/a/11/CPCSEA.).

Drug and Chemicals

The Methanolic extract of *Opuntia ficus indica* flowers was used as experimental extract. Phenytoin (Karnataka fine chem- Bangalore, India) used as standard antiepileptics agent. Pentylenetetrazole, (PTZ) (Karnataka fine chem- Bangalore), Strychnine (S D fine Chemicals – Mumbai) were all dissolved in normal saline. Dopamine, Nor-adrenaline (Neon Laboratory LTD – Mumbai), Adrenaline (Medlife Health Science LTD – Mumbai), Methanol (Nice Chemicals Private Limited, Kerala, India).

Preliminary phytochemical studies^[10-13]

Methanolic extract of *Opuntia ficus indica* was subjected to qualitative chemical tests for the identification of their active constituents. Test for the presence of carbohydrates, glycoside, resin, tannin, alkaloid, flavanoid, terpenoid, protein, saponin, anthraquinone, Coumaric, narcission, ascorbic acid, fibers, vitamins, iron, minerals, protein and amino acid were conducted as per the standard procedure. And the results obtained are presented in table 1.

Screening of Antiepileptic activity

Electroshock Induce Convulsion

Electro-convulsion shock, induce Hind Leg Tonic Extension (HLTE) in 99% of the animal. The MES test, developed by the Toman and Collaborators more than 60 years ago and probably the best validated test that predict drugs effective against generalized seizure of the tonic-clonic type. MES induce tonic extension has been suggested to identify drug that are effective in preventing seizure spread through neural tissue.^[14]

Mice weighting 20-30g received an electrical stimulus of 50 mA for 0.2 sec through auricular electrode to induce maximal seizure of their hind limbs, with tonic extension as the endpoint of the test.

The MES induced convulsion in animal represent grandmal type of epilepsy. This test can easily be conducted and requires a minimal investment in equipment and technical expertise, and is well standardized.^[15]

Experimental design

Animals were divided into 4 groups of 6 animals in each. Group I served as the control (0.9% saline, *p.o.*), Group II served as standard treated with phenytoin (25mg/kg, *i.p.*) and groups III and IV served as the low dose of MEOF (250mg/kg, *p.o.*) and high dose of MEOF (500mg/kg, *p.o.*) respectively. Treatment of MEOF to Group III and Group IV was given for 21 days. On 21st day after 30 minutes of *i.p.* injection of phenytoin and 60 minutes of oral administration of extract electric shock was given. Immediately after giving the electric shock, hind leg extension was observed. Decreased duration of hind leg extension was considered as a protective action and determined for each group.

Group	Treatment
Group I	Vehicle control (0.9% saline + electric stimulus(50mA;2 sec-duration)
Group II	Phenyton (25mg/kg, <i>i.p.</i>) + electric stimulus(50mA;2 sec-duration)
Group III	Test drug (250mg/kg, <i>p.o.</i>) + electric stimulus(50mA;2 sec-duration)
Group IV	Test drug (500mg/kg, <i>p.o.</i>) + electric stimulus(50mA;2 sec-duration)

Pentylenetetrazole Induce Convulsion

Pentylenetetrazole (PTZ) has been used widely to induce the convulsion since this chemical is highly sensitive for comparing different chemicals under standard conditions.^[16] It is considered as the model for petit-mal epilepsy. PTZ may elicit seizure by inhibiting the GABAergic mechanism.^[17,18] The dose of PTZ (80 mg/kg, *s.c.*) used in this study was determined and found to be approximately minimal dose (CD₁₀₀) that induces convulsion in 100% mice (i.e. control group). Mice that received PTZ were observed for 30 minutes.^[15]

Experimental design

Swiss albino mice of either sex were divided into 4 groups, each containing 6. Group I was considered as the vehicle control (0.9% saline, *p.o.*), group II was considered as the standard treated with Phenyton (25mg/kg, *i.p.*) and groups III and group IV were considered as the low dose of MEOF (250mg/kg, *p.o.*) and high dose of MEOF (500mg/kg, *p.o.*) respectively. On the 21st day of the experiment, after the administration of phenytoin and Methanolic extract of *Opuntia ficus indica* (MEOF), 30 minutes and 60 minutes respectively, PTZ injection 80mg/kg, *i.p.* was given. The animals were observed individually in the cage for an hour. The time interval between PTZ-injection and occurrence of seizures were measured. The delay of onset was calculated in comparison with the control group.

Group	Treatment
Group I	Vehicle control (0.9% saline + PTZ (80mg/kg, <i>i.p.</i>))
Group II	Phenyton (25mg/kg, <i>i.p.</i>) + PTZ (80mg/kg, <i>i.p.</i>)
Group III	Test drug (250mg/kg, <i>p.o.</i>) + PTZ (80mg/kg, <i>i.p.</i>)
Group IV	Test drug (500mg/kg, <i>p.o.</i>) + PTZ (80mg/kg, <i>i.p.</i>)

Strychnine induce convulsion

Experimental design

Swiss albino mice of either of sex were divided in the 4 group, each containing 6. Group I was consider as the vehicle control (0.9% saline), group II was consider as the standard treated with Phenyton (25mg/kg, *i.p.*) and group III and group IV was consider as the low dose of MEOF (250mg/kg, *p.o.*) and high dose of MEOF (500mg/kg, *p.o.*) respectively. On the 21st day of the experiment, after the administration of the phenytoin and Methanolic extract of *Opuntia ficus indica* (MEOF), 30 minutes and 60 minutes respectively, strychnine nitrate (2mg/kg, *i.p.*) was given. The time until occurrence of tonic extensor convulsion and death were noted during one hour period.

Group	Treatment
Group I	Vehicle control (0.9% saline + strychnine nitrate (2mg/kg, <i>i.p.</i>))
Group II	Phenyton (25mg/kg, <i>i.p.</i>) + strychnine nitrate (2mg/kg, <i>i.p.</i>)
Group III	Test drug (250mg/kg, <i>p.o.</i>) + strychnine nitrate (2mg/kg, <i>i.p.</i>)
Group IV	Test drug (500mg/kg, <i>p.o.</i>) + strychnine nitrate (2mg/kg, <i>i.p.</i>)

Brain amine estimation^[19]

Biogenic amine external standards including Norepinephrine, dopamine, were prepared as 0.5µM solution in 0.1N perchloric acid (PCA). For the preparation of tissue mice were decapitated and their brain was rapidly removed and dissected at 5°C. Brain sample were immediately homogenized in 20 vol (wt/vol) of ice-cold 0.1N perchloric acid (PCA) which contain 0.5µM dihydroxybenzylamine (DHBA); the homogenated was centrifuged for 20 min for 15,000 ×g at 2°C to precipitate protein. For measurement of dopamine in the striatum, the supernatant fluid was diluted 10- fold with 0.1N PCA to obtain concentration of endogenous dopamine in the linear range of the assay. Five to fifteen µl of PCA extract was injected into HPLC system.

To prepare the mobile phase 6.8 gram of sodium acetate, 100mg of disodium EDTA and 1 gram heptone sulfonate were diluted in 900ml of deionised and glass distilled water. After pH was adjusted to 4.0 with acetic acid the column adjusted to 935 ml, the solution was filtered

through a 0.45 μ M Millipore filter. To this solution, 65 ml of acetylnitrile was added; and buffer was de-gassed under vacuum.

HPLC was run at 254 nm and 280nm. Best peak were obtain at 280nm. The chromatograms were selected from the 280 nm.

Statistical Analysis

All the values were expressed as mean \pm SEM. Statistical analysis was performed by one way analysis of variance (ANOVA) followed by Tukey's posthoc tests. P values < 0.01 were considered as stastically significant.

RESULTS

Table 1: Effect of Methanolic extract of *Opuntia ficus indica* flowers (MEOF) on Maximal electroconvulsive shock (MES) induced seizures in mice.

Group	Treatment	Flexion (sec)	Extensor (sec)	Clonus (sec)	Stupor (sec)	% protection against extensor	% protection against mortality
I	Vehicle control (0.9% saline, <i>p.o.</i>)	2.317±0.1078	19.00 ±1.592	11.50±0.7638	17.75±3.435	0	50
II	MEOF (250mg/kg, <i>p.o.</i>)	1.783±0.1138**	13.33±1.022**	9.417±0.5974*	11.03±1.415*	29.84	66.66
III	MEOF (500mg/kg, <i>p.o.</i>)	1.133±0.1054***	8.500±0.6708***	5.433±0.1978***	7.450±0.2527**	55.26	100
IV	Phenytoin (25mg/kg, <i>i.p.</i>)	0.7000±0.09661***	1.833±0.7177***	3.783±0.1046***	4.633±0.1783***	90.35	100

Values are expressed as mean ± SEM (n=6).

Statistical significant test for comparison was done by ANOVA, followed by Tukey's posthoc Test.

*p<0.05, **p<0.01 and ***p<0.001 as compared to control group.




Figure 1: Effect of Methanolic extract of *Opuntia ficus indica* flowers (MEOF) on MES induced tonic seizure in mice.

HLTE = Hind Leg Tonic Extension

Each bar represent mean±SEM (n=6)

Statistical significant test for comparison was done by ANOVA, followed by Tukey's posthoc Test.

*p<0.05 and ***p<0.001 as compared to the control group.

Table 2: Effect of ethanolic extract of *Opuntia ficus indica* flowers (MEOF) on PTZ induced epilepsy in mice.

Group	Treatment	Onset of convulsion (sec)	% protection against convulsion	% mortality
I	Vehicle control (0.9% saline, <i>p.o.</i>)	155.8±8.101	0	100
II	MEOF (250mg/kg, <i>p.o.</i>)	207.3±7.531**	33.05	66.66
III	MEOF (500mg/kg, <i>p.o.</i>)	236.3±11.14***	51.66	50
IV	Phenytoin (25mg/kg, <i>i.p.</i>)	292.5±15.15***	87.74	33.33

Values are expressed as mean ± SEM (n=6).

Statistical significant test for comparison was done by ANOVA, followed by Tukey's posthoc Test.

*p<0.05, **p<0.01 and ***p<0.001 as compared to control group.




Figure 2: Effect of ethanolic extract of *Opuntia ficus indica* flowers (MEOF) on PTZ induced epilepsy in mice.

Each bar represent mean \pm SEM (n=6)

Statistical significant test for comparison was done by ANOVA, followed by Tukey's posthoc Test.

*p<0.05, **p<0.01 and ***p<0.001 as compared to the control group.

Table 3: Effect of Methanolic extract of *Opuntia ficus indica* flowers (MEOF) on Strychnine induced epilepsy in mice.

Group	Treatment	Onset of convulsion (sec)	% protection against convulsion	% mortality
I	Vehicle control (0.9% saline, <i>p.o.</i>)	154.2 \pm 7.236	0	100
II	MEOF (250mg/kg, <i>p.o.</i>)	195.0 \pm 8.383**	26.45	50
III	MEOF (500mg/kg, <i>p.o.</i>)	230.8 \pm 5.231***	49.67	33.33
IV	Phenytoin (25mg/kg, <i>i.p.</i>)	290.0 \pm 12.38***	88.06	16.66

Values are expressed as mean \pm SEM (n=6).

Statistical significant test for comparison was done by ANOVA, followed by Tukey's posthoc Test.

***p<0.001 as compared to control group.




Figure 3: Effect of Methanolic extract of *Opuntia ficus indica* flowers (MEOF) on Strychnine induced epilepsy in mice.


Each bar represent mean \pm SEM (n=6).

Statistical significant test for comparison was done by ANOVA, followed by Tukey's posthoc Test.

***p<0.001 as compared to the control group.

Table 4: Effect of MEOF on neurotransmitter levels in mice brain after MES induced epilepsy.

Group	Design of treatment	Concentration of Dopamine ($\mu\text{g}/\text{gm}$)	Concentration of Noradrenaline ($\mu\text{g}/\text{gm}$)
I	Vehicle control (0.9% saline, <i>p.o.</i>)	0.02354	0.0491
II	MEOF (250mg/kg, <i>p.o.</i>)	0.04929	0.0703
III	MEOF (500mg/kg, <i>p.o.</i>)	0.10371	0.1135
IV	Phenytoin (25mg/kg, <i>i.p.</i>)	0.1548	0.1712




(1)




(2)


Figure 4.1: Chromatogram of Standard Dopamine (1) and Standard Noradrenaline (2)




(1)



(2)



(3)



(4)

Figure 4.2: MEOF chromatogram of control (1) low dose (2) high dose (3) standard (4) for Dopamine and Noradrenaline.




Figure 4.3: Effect of Methanolic extract of *Opuntia ficus indica* flowers (MEOF) on brain amine (Dopamine) on MES induced epilepsy in mice.




Figure 4.4: Effect of Methanolic extract of *Opuntia ficus indica* flowers (MEOF) on brain amine (Nor-adrenaline) on MES induced epilepsy in mice.

Table 5: Phytoconstituents present in 65% Methanolic extract of *Opuntia ficus indica* flowers.

Serial no.	Phytoconstituents	Result
1	Carbohydrates	Present
2	Alkaloids	Present
3	Saponins	Present
4	Flavonoids	Present
5	Phenolics	Present
6	Total Flavonoids	Present
7	Glycosides	Present
8	Proteins	Present
9	Amino acids	Present
10	Phytosterol	Present
11	Tannins	Present
12	Terpenoids	Present
13	Polypherol	Present

DISCUSSION

Epilepsy is a chronic neurological condition, characterized by recurrent seizures that are caused by abnormal cerebral nerve cell activity. The treatment of epilepsy is always a challenge for researchers and clinical practitioners. However, several new drugs have been introduced for the treatment of epilepsy. Despite this progress, about 30% of patients with epilepsy are resistant to current pharmacotherapies and many of the available antiepileptic drugs. Based on this reason, there has been a continuous attempt to find new antiepileptic drugs which possessing multiple mechanisms of action with lesser adverse effect.^[20]

MES induced seizures in the animal is most frequently used model for the identification of anticonvulsant activity of the drugs against generalized tonic-clonic seizure “grand mal” and coritical focal seizures. PTZ is a Chemoconvulsants, which induces seizures by the inhibition of GABA-A receptors and it is widely accepted experimental model for “generalized absence seizure”^[21-23] and also a valid model for human generalized myoclonic seizures and generalized seizures of the petitmal type.

MES induced tonic extension can be prevented by AEDs that inhibit either voltage gated channels or by blocking glutamergic excitation mediated by the N-methyl-D-aspartate (NMDA) receptor.^[24] AEDs have the ability to open K⁺ channels in neuronal cells; can prevent seizure induced by PTZ and MES. Currently used anticonvulsant drugs (e.g. phenytoin, carbamazepines) effective in therapy of generalized tonic-clonic and partial seizures have been found to show strong anticonvulsant action in MES test.^[25-28]

The treatment with MEOF on mice significantly reduced tonic hind leg extension in MES induced epilepsy. Since, MEOF significantly inhibited generalized tonic-clonic seizures in MES test; which may be either inhibiting voltage gated channel or blocking glutamergic transmission.

PTZ is non-competitive antagonist which blocks GABA mediated Cl^- influx and leads to enhanced release of excitatory neurons to release NTs such as α -amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid receptor (AMPA) and N-methyl-D-aspartic acid (NMDA). This results in neuronal excitotoxicity leading to seizures and convulsions characterized by clonic and tonic phases.^[29] so the NMDA receptor antagonist ketamine, diazocilpine have reported the effect against PTZ induced seizure.^[30]

The antagonism of PTZ-induced seizures suggests the interaction of the Methanolic extract *Opuntia ficus indica flowers* either with the potentiating of GABAergic neurotransmission system and/ or inhibition of NMDA receptors.

Strychnine may cause seizures by activation of excitatory amino acid NMDA receptor. Since Strychnine has been shown to interact with the NMDA neurotransmission thus Strychnine-induced seizures can be prevented by drugs that antagonize NMDA-receptor-mediated excitatory neurotransmission.^[31]

There are five established biogenic amine neurotransmitters, of which there are three catecholamines Dopamine, Norepinephrine (Nor-adrenaline) and Epinephrine (Adrenaline), and other biogenic amines are histamine and serotonin.^[32]

The brain amine estimation was done by the HPLC method in which increase of amine level in the forebrain region of treated mice shows the significant antiepileptic activity. Increase of brain monoamine level by inhibiting the monoamine oxidase (MAO), an enzyme responsible for the seizure by destruction of amines in brain. Inhibition of prostaglandin synthesis increases the levels of dopamine and nor-adrenaline, which also cause an inhibition of seizure activity.^[33,34]

In the present study the administration of MEOF significantly increased the brain levels of dopamine and nor-adrenaline attributed to the significant protection against MES induced seizures (Table 5).

CONCLUSION

The present study suggests that the Methanolic extract of *Opuntia ficus indica* flowers has significant antiepileptic activity against various models of epilepsy in experimental mice. This activity of *Opuntia ficus indica* flowers might be due to presence of flavonoids (rutin & quercetin active principles of extract). Further studies and tests are required to evaluate the exact mechanisms, specific active principles and the safety profile of the plant as a medicinal remedy for convulsion.

ACKNOWLEDGEMENT

Please refer some articles and write.

Example: I thank Principal Dr. Shiva kumar Swamy for providing facilities to carry out the research at Mallige college of Pharmacy, Sivepura Bangalore.

REFERENCES

1. Singh P, Mishra G, Garg VK, Jha KK, Khosa RL. Phytochemical screening and anticonvulsant activity of alcoholic extract of root of vitex negundo. *International journal of pharmaceutical research and innovation*, 2011; 4: 16-9.
2. Mattson RH. Efficacy and adverse effects of established and new antiepileptic drugs. *Epilepsia*. 1995; 36(2): 13-26.
3. Nieoczym D, Socala K, Raszewski G, Wlaz P. Effect of quercetin and rutin in some acute seizure models in mice. *Progress in Neuro-Psychopharmacology & Biological Psychiatry*. 2014; 54: 50-8.
4. Goyal RK, Mehta AA, Balraman R, Burande MD. Derasari and Gandhi's elements of Pharmacology. 16th ed. Ahmedabad(India): B S Shah prakashan; 2007; 233-6 p.
5. Debnath S, Kannadasan M, Ghosh S, Ghosh NS, Chakraborty R, Sen S. Antiepileptic activity of the hydroalcoholic extract of erythrina fusca lour bark against the animal model of MES, PTX and PTZ induced epileptic seizure models. *international journal of chemistry research*. 2010; 1(2): 3-5.
6. Dwivedi C. Antiepileptic Drugs. *Am J Phar E*, 2001; 65: 197- 202.
7. Rang HP, Dale MM, Ritter JM, Flower RJ. Rang and Dales's Pharmacology. 6th ed. New York: Churchill Livingstone, Edinburgh: Elsevier; 2007; 575-84 p.
8. Aliyu MM, Musa AI, Kamal MJ, Mohammed MG. Photochemical screening and anticonvulsant studies of ethyl acetate fraction of *Globimetula braunii* on laboratory animals. *Asian Pacific Journal of Tropical Biomedicine*. 2014; 4(4): 285-9.

9. Loscher W. Critical review of current animal models of seizures and epilepsy used in the discovery and development of new antiepileptic drugs. *Seizure european journal of epilepsy*. 2011; 20(5): 359-68.
10. De Leo M, Abreu MBD, Pawlowska AM, Cioni PL, Braca A. Profiling the chemical content of *Opuntia ficus-indica* flowers by HPLC–PDA-ESI-MS and GC/EIMS analyses. *Phytochemistry Letters*. 2010; 3(1): 48-52.
11. Yeddes N, Cherif JK, Guyot S, Sotin H, Ayadi MT. Comparative Study of Antioxidant Power, Polyphenols, Flavonoids and Betacyanins of the Peel and Pulp of Three Tunisian *Opuntia* Forms. *antioxidants*. 2013; 2: 37-51.
12. Alimi H, Hfaiedh N, Bouoni Z, Sakly M, Rhouma KB. Evaluation of antioxidant and antiulcerogenic activities of *Opuntia ficus indica* f. *inermis* flowers extract in rats. *Environmental Toxicology and Pharmacology*. 2011; 32(3): 406-16.
13. Rabhi A, Limam F, Fallech H, Ksouri R, Abdelly C, Raies A. *Opuntia ficus indica* flower: A prominent source of phenolic compounds and unsaturated fatty acid. *International journal of advanced research*. 2015; 3(4): 41-55.
14. Branci CMM, Alves GL, Figueiredo IV, Falcao AC, Caramona MM. The maximal electroshock seizure (mes) model in the preclinical assessment of potential new antiepileptic drugs. *Methods Find Exp Clin Pharmacol*. 2009; 31(2): 101-6.
15. Raza M, Alghasham AA, Alorainy MS, El-Hadiyah TM. Beneficial Interaction of Thymoquinone and Sodium Valproate in Experimental Models of Epilepsy. *Reduction in Hepatotoxicity of Valpro Sci Pharm*. 2006; 74: 159-73.
16. Mairhadi K, Daruush M, Reza A. Effect of Lovastatin on PTZ-Induced Seizure Threshold in Mice. *Global Veterinaria*. 2011; 7(4): 386-90.
17. Hegde K, Thakker SP, Joshi AB, Shastry CS, Chandrashekhar KS. Anticonvulsant activity of *Carissa carandas* Linn. root extract in experimental mice. *Tropical Journal of Pharmaceutical Research*. 2009; 8(2): 117-25.
18. Forsgren L, Beghi E, Oun A, Sillanpaa M. The epidemiology of epilepsy in Europe: a systematic review. *Eur J Neurol*. 2005; 12: 245-53.
19. Zaczek R, Coyle JT. Rapid and simple method for measuring biogenic amines and metabolites in brain homogenates by HPLC electrochemical detection. *J Neural Transmission*. 1982; 53: 1-5.
20. Suong LK, Phuong Lan TT, Nguyen VP. Anti-convulsive activity of NL197, A derivative from 4(3H)Quinazolinon on chemical induced seizure mice. *13rd International Electronic Conference on Synthetic Organic Chemistry (ECSOC)*. 2009; 2-30.

21. Loscher W, Schmidt D. Which animal models should be used in the search for new antiepileptic drugs? A proposal based on experimental and clinical considerations. *Epilepsy Res.* 1988; 2(3): 145-81.
22. Oliveira FA, Almeida RN, Sousa MFV, Barbosa-Filho JM, Diniz SA, Medeiros IA. Anticonvulsant properties of N-salicyloyltryptamine in mice. *Pharmacol Biochem Behav.* 2001; 68: 199-202.
23. Singh P, Garg VK, Sharma PK, Gupta S. Antiepileptic activity of aqueous extract of *Tricosanthes dioica* Roxb. *Asian Journal of Plant Science and Research.* 2012; 2(1): 45-7.
24. Subramaniam S, Rho JM, Penix L, Donevan SD, Fieldin RP, Rogawski MA. Felbamate block the N-methyl-Daspartame receptor. *J Pharmacol Exp Ther.* 1995; 273: 878-86.
25. Gopi G, Jayasri P, Elumalai A. Anti-epileptic activity of malachra capitata L. on maximal electroshock (MES) and pentylenetetrazole (PTZ) induced seizures models. *International Journal of Pharmacology & Toxicology.* 2012; 2(2): 104-8.
26. Kupferberg HJ. Antiepileptic drug development program: a cooperative effort of government and industry. *Epilepsia.* 1989; 30(1): 51-6.
27. Stables JP, Kupferberg HJ, Levy RH, Mattson RH, Meldrum BS. The NIH Anticonvulsant Drug Development (ADD) Program: Preclinical Anticonvulsant Screening project. In: *Antiepileptic Drugs.* 1995; 4: 4-17.
28. Patten D, Foxon GR, Martin KF, Hallwell RF. An electrophysiological study of the effects of propofol on native neuronal ligand-gated ion channels. *Clin Exp Pharmacol Physiol.* 2001; 28: 451-8.
29. Eggadi V, Kulandaivelu U, Sharvanabhava BS, Jupally VR. Screening of the anticonvulsant activity of some isatin derivatives in experimental seizure models and Its effect on brain GABA levels in mice. *American Journal of Pharmacological Sciences.* 2013; 1(3): 42-6.
30. Sundriyal A, Brijjem KRV, Kalia A. Antiepileptic potential of anisomeles indica (Linn.) kuntze aerial parts in pentylenetetrazole-induced experimental convulsion in wister rats. *Indian Journal of Experimental Biology.* 2013; 51: 715-20.
31. Larson AA, Beitz AJ. Glycine Potentiates Strychnine-Induced Convulsions: Role of NMDA Receptors. *The Journal of Neuroscience.* 1988; 8(10): 3822-6.
32. Walker R, Whittlesea C. *Clinical Pharmacy and Therapeutics.* 5th ed. Edinburgh: Churchill Livingstone/Elsevier; 2012; Xii: 983 p. p.

33. Rajasree PH, Singh R, Sankar C. Screening for antiepileptic activity of Moringa oleifa root extract. *Int J of Pharm & Life Sci (IJPLS)*. 2012; 3(11): 2115-9.
34. Muralidharan P, Srikanth J. Antiepileptic activity of Morinda citrifolia Linn fruit extract. *E-Journal of chemistry*. 2010; 7(2): 612-6.