

**MOLECULAR STUDY OF FOUR SPECIES OF *SARCOCYSTIS*
ISOLATED FROM SHEEPS ESOPHAGUS AND INTER SKELETON
MUSCLE IN BAGHDAD.**

Safa T. Whaeb^{*}, Azhar A. Faraj and Ilham A. Khalaf

*Veterinary Medicine of Parasitology DEpartmnt for Post Graduate Studies, Baghdad
University.

Al-Razi Center for Research and Diagnostic Kits, Ministry of Industry and Minerals,
Baghdad, Iraq.

Article Received on
06 June 2016,

Revised on 27 June 2016,
Accepted on 17 July 2016

DOI: 10.20959/wjpr20168-6654

***Corresponding Author**

Dr. Safa T. Whaeb

Veterinary Medicine of
Parasitology DEpartmnt
for Post Graduate Studies,
Baghdad University.

ABSTRACT

Sarcocystis is an obligatory intracellular protozoan parasite which can infect humans and animals. Sheep are intermediate hosts of four *Sarcocystis* species: *Sarcocystis tenella*, *Sarcocystis gigantea*, *Sarcocystis arieticanis* and *Sarcocystis medusififormis*. The purpose of this study was to perform a molecular identification of the macroscopic and microscopic cysts of *Sarcocystis* in sheep. In this investigation, the microscopic cysts of *Sarcocystis* were assessed in slaughtered sheep. The digestion method was used for bradyzoites observation in, esophagus and inter costal muscle samples ($P < 0.01$). PCR analysis was conducted on microscopic cysts and also all other samples. Sequencing was performed for ten PCR products. Genotypes were identified by

BLAST search and homology analysis. Digestion method and PCR analysis revealed positive results in all samples taken from esophagus and inter costal muscle. Genotyping of 5 tissue samples proved that the genotype of macroscopic belonged to microscopic cysts to *Sarcocystis tenella*. Microscopic cysts are more prevalent than macroscopic cysts and they can cause enormous economic losses.

KEYWORDS: *sarcocystis* spp, polymerase chain reaction (PCR), Iraq.

INTRODUCTION

species are intracellular protozoan parasites infecting a wide range of livestock. Some of *Sarcocystis* genus are pathogenic for animals such as sheep and cattle which cause enormous economic losses.^[1] Studies in different regions of the world indicate that the prevalence of *Sarcocystis* infection in slaughtered cattle and sheep are between 70% to 100%^[2,3] Additionally, studies in Iran showed that the prevalence of this parasite in the animal was between 85% to 100%^[4,5]. For example, studies in Kerman and Ahwaz provinces indicated that 100% of animals were infected with *Sarcocystis*.^[5, 6] Different species of *Sarcocystis* have been isolated from animals worldwide. *Sarcocystis tenella* was isolated from sheep in Iran and Brazil.^[7, 8] In another study, *Sarcocystis moulei* was reported from reindeer.^[9] Also, Nourani et al. isolated *Sarcocystis hominis* from cattle^[10] while Kalantari et al. separated *S. cruzi* from cattle^[11] determined *S. gigantea* and *S. arieticanis* in sheep.^[12]

DIAGNOSIS

The diagnosis is usually made *post mortem* by examination of the skeletal muscle. In some species the cysts may be visible to the naked eye (ducks, mice, rabbits and sheep) but in most microscopic examination is required. *Ante mortem* diagnosis may be made with the use of dermal sensitivity testing or complement fixation tests. Muscle biopsy is also diagnostic but this is much less commonly used.^[13]

Oocysts with two sporocysts or individual sporocysts in human feces are diagnostic of intestinal infection.

The conventional tools for species diagnosis of *Sarcocystis* spp. were based on transmission electron microscopy, structure of the cyst wall in the striated muscles of the intermediate host or information about the lifecycle of the parasite.^[14] However, because of showing the morphologic variations in these procedures they are not exactly reliable at the species-specific identification. On the other hand, electron microscopy is not a choice for wide and extensive detective studies.^[13]

In recent times, various molecular techniques such as PCR and its variants based on sequence changes have been used regarding the sensitivity and rapidity to determine genetic diversity among many parasites, phylogenetic and taxonomic studies and in epidemiological mapping.^[15]

Thus, definitive diagnosis of sarcocystosis requires identification of sporocysts in feces. However, the sporocysts of different species are similar in size and shape, making species identification almost impossible by microscopy. Therefore, sequencing of the small subunit ribosomal RNA (18S rRNA) gene was introduced as an ideal means for species-specific detection. In fact, this gene contains hypervariable regions interspersed within highly conserved DNA sequences, making it ideal for differentiation between species.

MATERIALS AND METHODS

Sarcocystis spp. strains were isolated from organs samples (esophagus and skeleton muscle) from 60 sheep selected randomly from Butcher at different localities of Baghdad, Iraq between February 2016 to July 2016. Isolation and identification of the strains were made by conventional methods^[13] conventional methods.

DIGESTION METHOD

Tissue digestion method was used for observing bradyzoites in the organ samples. Seventy grams of each tissue were ground and digested in 1.5% HCL acid and 0.5% pepsin at 29 °C overnight. The digested samples were filtered through mesh and centrifuged 1500 RPM for 10 min. Then, the supernatant fluid was discarded and sediment was stained with Gimsa and examined microscopically for detecting bradyzoites of *Sarcocystis*.

PROCESSING OF THE SAMPLES FOR PCR ASSAY

A volume of 1.5 ml of the post-enriched sample was centrifuged at 14,000g for 1 min, DNA was extracted using Presto Mini g DNA Tissue Kit according to manufacturer's instructions (Geneaid, Korea). The extracted DNA was stored -20 °C until use. The extracted DNA then quantified through measurement of its OD₂₆₀ by ND-2000 spectrophotometer (Thermo Scientific Inc., USA).

PCR AMPLIFICATION ANALYSIS

The virulence determinants investigated using the oligonucleotide primers included the gene *SARI*. For all the gene, The polymerase chain reaction (PCR) amplification was performed in a final volume of 20µl containing 10 Intron- Master Mix (KOBA) which contains (Taq polymerase, PCR buffer, MgCl₂ and dNTPs), 200 ng of DNA template added 1µl of 10 pmol each primer, and 6µl of nuclease free water, in the present study, the amplification parameters and primer sequence were used in (table1). The amplification of gene was carried out with Master cycler (Eppendorf, Germany). Amplified products were separated by agarose gel

electrophoresis (2% agarose containing 0.5 mg ethidium bromide in 0.5 × Tris - EDTA electrophoresis buffer) at 5V/cm for 2h and photographed under UV illumination.

RESULTS AND DISCUSSIONS

In this study, 60 samples of muscle and esophagus had Microscopic cysts (table1). In addition, these organs were found to be infected with microscopic cysts by digestion method. The results of digestion method showed that all samples of muscle and esophagus were infected with bradyzoites of *Sarcocystis*. PCR analysis of microcysts as well as all samples showed a specific 600 bp band on the agarose gel (Figures 2). The results obtained from sequencing of five samples (3 muscles and 2 esophagus). showed that the genotype of microscopic cysts to *Sarcocystis tenella*.

Microscopic examination by trichinoscopy Technique direct between two slid for appear diameter (13-1)*(7-0.5) like figure (1).



Figure (1): show one types of *sarcocystis* in sheep in different organ X40

Table 1. Number of microscopic cysts in sheep's tissues and used Methods.

organ	Microscopic cyst	Digestion method	Molecular method	negative
esophagus	40	40	40	0
muscles	20	20	20	0




Figure (2): Gel electrophoresis of 1% agarose gel stained with ethidium bromide for DNA extraction of *Sarcocystis*.

DISCUSSION

All samples (muscle and esophagus) were infected with *Sarcocystis* spp. The studies in Baghdad and other parts of the world indicated that live stocks are infected with *Sarcocystis* spp.^[16] Other studies in different provinces of Iran showed that 97% of sheep had *Sarcocystis* infection.^[17] In previous studies throughout the world, species of *Sarcocystis* were isolated from different animals.^[18, 19] DaSilva and Langoni isolated *S. tenella* from the sheep in Brazil.^[18] Al- Hoost et al. reported *S. moulei* from the sheep in Saudi Arabia.^[20] Moreover, Gjerde isolated and characterized *S. grueneri* from reindeer based on molecular method.^[19] In other studies in Iran, *Sarcocystis hominis* and *Sarcocystis cruzi* were studied in cattle.^[21] isolated *S.gigantea* and *S. arieticanis* from the sheep by PCR-RFLP method in Qazvin province, Iran.^[6] Furthermore, other researchers reported *S miescheriana* from boar^[21] and *S. tenella* from sheep in Iran.^[10] Additionally, Mahran in Egypt using morphometric method indicated that *S gigantea* and *S. tenella* caused macroscopic and microscopic cysts.^[22] Using daub smear method, showed that 91% of cows were infected with microscopic cyst and did not have any macroscopic cysts.^[23] Kargar Jahromi et al. using digestion method proved that goats had microscopic and macroscopic cysts.^[24] Molecular analysis was not performed in the above studies, but, the use of molecular methods in the present study showed that *S. gigantea* and *S. tenella* can cause macroscopic and microscopic cysts, respectively. The results of this investigation was in accordance with Anja and Astrid's study who reported that *S. gigantea* and *S. tenella* can cause macroscopic and microscopic cysts, respectively^[10] *S.*

tenella is among the pathogenic species and can induce microscopic cysts. The severity of clinical symptoms caused by this species depends on the dose of ingested sporocysts and the immune system of the host.^[24,25] *S. tenella* can lead to acute sarcocystosis in uninfected sheep.^[7] Nonspecific infection symptoms include fever, anorexia, tachycardia and anemia could be observed following infection. In acute sarcocystosis, central nervous system will be involved, and it can cause encephalitis and encephalomyelitis and subsequently death in sheep.^[26,27] In pregnant sheep, acute sarcocystosis can cause fetal death or premature birth of offspring. Chronic sarcocystosis can create economic problems due to reduced meat, milk and wool.^[27,28,29] Also, Dubey reported that *S. tenella* caused symptoms such as inflammation, hepatitis and myocarditis in sheep inoculated with *S. tenella* sporocysts from canine feces.^[30]

REFERENCE

1. Heckerth AR, Tenter AM. Comparison of immunological and molecular methods for the diagnosis of infections with pathogenic Sarcocystis species in sheep. Tokai J Exp Clin Med., 1998; 23: 293-302.
2. Pereira A, Bermejo M. Prevalence of Sarcocystis cysts in pigs and sheep in Spain. Vet Parasitol., 1988; 27: 353-5.
3. Woldemeskel M, Gebreab F. Prevalence of sarcocysts in livestock of northwest Ethiopia. Zentralbl Veterinarmed B., 1996; 43: 55-8.
4. Oryan A, Ahmadi N, Mousavi SM. Prevalence, biology, and distribution pattern of Sarcocystis infection in water buffalo (*Bubalus bubalis*) in Iran. Trop Anim Health Prod., 2010; 42: 1513-8.
5. Hamidinejat H, Razi Jalali MH, Nabavi L. Survey on Sarcocystis Infection in Slaughtered Cattle in South-West of Iran, Emphasized on Evaluation of Muscle Squash in Comparison with Digestion Method. J Animal Vet Advances., 2010; 9: 1724-6.
6. Nourollahi Fard SR, Asghari M, Nouri F. Survey of Sarcocystis infection in slaughtered cattle in Kerman, Iran. Trop Anim Health Prod., 2009; 41: 1633-6.
7. da Silva RC, Su C, Langoni H. First identification of Sarcocystis tenella (Railliet, 1886) Moule, 1886 (Protozoa: Apicomplexa) by PCR in naturally infected sheep from Brazil. Vet Parasitol., 2009; 165: 332-6.
8. Shahbazi A, Falah S, Khanmohamadi M, et al. Identification of Sarcocystis tenella and Sarcocystis articanis from slaughtered sheep using PCR-RFLP in Tabriz province. Journal of pathobiology., 2013; 10: 959-64. [In Persian]

9. Gjerde M. Ultrastructure of the cysts of *Sarcocystis grueneri* from cardiac muscle of reindeer (*Rangifertarandustarandus*). *Z Parasitenkd.*, 1985; 71: 189-98.
10. Nourani H, Matin S, Nouri A, et al. Prevalence of thin-walled *Sarcocystis cruzi* and thick-walled *Sarcocystis hirsuta* or *Sarcocystis hominis* from cattle in Iran. *Trop Anim Health Prod.*, 2010; 42: 1225-7.
11. Kalantari N, Bayani M, Ghaffari S. *Sarcocystis cruzi*: First molecular identification from cattle in Iran. *Int J Mol Cell Med.*, 2013; 2: 125-30.
12. Dalimi Asl AH, Mutamedi G, Paykari H, et al. Detection of *Sarcocystis* spp. of slaughtered sheep in Gazvin Ziaran slaughter house by molecular assay. *Journal of modarres Medical Science.*, 2008; 11: 65-72. [In Persian]
13. Van den Enden, E. M. Praet, R. Joos, A. Van Gompel, and P. Gigasse. Eosinophilic myositis resulting from sarcocystosis. *J. Trop. Med. Hyg.*, 1995; 98: 273-276.
14. Gasbarre L.C. Suter P. payer R. Humoral and cellular immune responses in cattle and sheep inoculated with *Sarcoc* B&dqo;. *B/is, Am. J. Vet. Res.* 8(1984) L!92-)596.
15. La Perle K.M., Silveria F., Anderson D.E., Blomme E.A. Dalmeny disease in an alpaca (*Lama pacos*): sarcocystosis, eosinophilic myositis and abortion. *J. Comp. Pathol.*, 1999; 121(3): 287-293.
16. Hamidinejat H, Hekmatimoghaddam SH, Jafari H, et al. Bahari P et al. Int J Mol Cell Med Winter 2014; 3-1. 56 Prevalence and distribution patterns of *Sarcocystis* in camels (*Camelus dromedarius*) in Yazd province, Iran. *J parasit Dis.*, 2013; 37: 163-65.
17. Shekarforoush SS, Shakerian A, Hasanpoor MM. Prevalence of *Sarcocystis* in slaughtered one-humped camels (*Camelus dromedarius*) in Iran. *Trop Anim Health Prod.*, 2006; 38: 301-3.
18. da Silva RC, Su C, Langoni H. First identification of *Sarcocystis tenella* (Railliet, 1886) Moule, 1886 (Protozoa: Apicomplexa) by PCR in naturally infected sheep from Brazil. *Vet Parasitol.*, 2009; 165: 332-6.
19. Al-Hoot AS, Al-Qureishy SA, Al-Rashid K, et al. Microscopic study on *Sarcocystis moulei* from sheep and goats in Saudi Arabia. *J Egypt Soc Parasitol.*, 2005; 35: 295-312.
20. Gjerde M. Ultrastructure of the cysts of *Sarcocystis grueneri* from cardiac muscle of reindeer (*Rangifertarandustarandus*). *Z Parasitenkd.*, 1985; 71: 189-98.
21. Kia EB, Mirhendi H, Rezaeian M, et al. First molecular identification of *Sarcocystis miescheriana* (Protozoa, Apicomplexa) from wild boar (*Sus scrofa*) in Iran. *Exp Parasitol.*, 2011; 127: 724-6.

22. Mahran OM. Sarcocystis infection in sheep and goats slaughtered in Shalatin Abattoir, Red Sea Governorate, Egypt. *Assiut Veterinary Medical Journal.*, 2009; 55: 341-55.
23. Bonyadian M, Meshki B. Study on infection of cow carcasses to Sarcocyst Spp in slaughtered cows in Shahrei. *Pajouhesh va sazandegi.*, 2006; 19: 14-8. [In Persian]
24. Kargar Jahromi Z, Solhjoo K, Zareian Jahromi M, et al. Investigation of Sarcocystis Infection in Slaughtered Goats in Jahrom Abattoir. *JFUMS.*, 2012; 2: 163-7.
25. O' Donoghue P, Rommel M. Australian-German collaborative studies on the immunology of Sarcocystis infections. *Angew Parasitol.*, 1992; 33: 102-19.
26. Ugglia A, Buxton D. Immune responses against Toxoplasma and Sarcocystis infections in ruminants: diagnosis and prospects for vaccination. *Rev Sci Tech.*, 1990; 9: 441-62.
27. Jeffrey M. Sarcocystosis of sheep., 1993; 15: 2-8. *In Practice.*, 1993; 15: 2-8.
28. Leek RG, Fayer R, Johnson AJ. Sheep experimentally infected with sarcocystis from dogs. I. Disease in young lambs. *J Parasitol.*, 1977; 63: 642-50.
29. Munday BL. The effect of Sarcocystis tenella on wool growth in sheep. *Vet Parasitol.*, 1984; 15: 91-4.
30. Collins GH, Charleston WA, Moriarty KM. Sarcocystis species in sheep. *N Z Vet.*, J 1976; 24: 123-4.
31. Dubey JP. Lesions in sheep inoculated with Sarcocystis tenella sporocysts from canine feces. *Vet Parasitol.*, 1988; 26: 237-52.