

## EXTRACTION AND CHEMICAL TESTS ON NIGELLA SATIVA L. COLLECTED FROM VIDARBHA REGION OF INDIA

Devender Sharma\*<sup>1</sup>, Kajal V. Kosankar<sup>2</sup> and Ameya M. Lanjewar<sup>3</sup>

Department of Pharmaceutics,

<sup>1</sup>Hi – Tech college of Pharmacy, Chandrapur (M.S.) India.

<sup>2</sup>MVP College of Pharmacy, Nashik Maharashtra (India).

<sup>3</sup>Kamla Nehru College of Pharmacy, Butibori, Nagpur Maharashtra (India).

Article Received on  
04 Dec. 2017,  
Revised on 25 Dec. 2017,  
Accepted on 14 Jan. 2018  
DOI: 10.20959/wjpr20182-10813

### \*Corresponding Author

Devender Sharma

Department of  
Pharmaceutics, Hi – Tech  
college of Pharmacy,  
Chandrapur (M.S.) India.

### ABSTRACT

*Nigella sativa* Linn (family- Ranunculaceae) is a widely used medicinal plant; especially in India this plant is largely used and popular in various Indigenous System of Medicine like Ayurveda, Siddha, Unani and Tibb. *Nigella sativa* is a wonder and very useful plant which has the potential to cure almost all diseases except death. The medicinal value of the plant has been reported quite long ago and finds its place in Holy Quran. *Nigella* seed oil is becoming popular in and out of the Islamic world. Composition *Nigella Sativa* of Family Ranunculaceae contains chemical components like Thymoquinone, alpha hederin which are anticancer in nature. Seeds of *Nigella sativa* L.

(black cumin or black seeds) are widely used in traditional Islamic medicine and for culinary purposes worldwide. So in this article emphasis on the different solvent extraction and its chemical tests have been shown from the seeds collected from Chandrapur, i.e. Vidarbha region of India. The seeds of this medicinal plant are used as astringent, stimulant diuretic and anthelmintic. These are also useful in jaundice, intermittent fever, paralysis and piles.

**KEYWORDS:** *Nigella sativa*, Medicinal plant, Thymoquinone, Anticancer, Seeds.

### INTRODUCTION

Medicinal plants are used for curing diseases for several centuries in numerous indigenous systems of drugs also as folk medicines. Natural products are awesome success in our society. The utilization of plant and microorganism secondary metabolites has aided in

doubling of our life within the twentieth century. Most of these activities have already been attributed to thymoquinone, a major component of the essential oil of the seeds (Ali and Blunden, 2003). Additionally, in its native range and far beyond, *N. sativa* seeds are also frequently used as spice and condiment in various recipes due to their characteristic aroma and bitter and peppery taste (Hedrick and Sturtevant, 1972) Finally, *N. sativa* seeds are used to prepare a highly prized nutritive oil. They need reduced pain and suffering, and revolutionized medication by enabling the transplantation of organs. Since their chemical diversity is predicated on biological and geographical diversity, the complete globe is explored for bioprospecting by researchers (Khare 2004). Cancer is one among the foremost threats of recent life, that is taken into account because the second reason behind death when myocardial infarction. Variant individuals die per annum in numerous types of cancer despite tremendous efforts to search out strategies of management and cure within the last century, nice advances were created in trendy bioscience to regulate malady (Sharma, Yelne et al. 2000). However several diseases like cancers aren't curable absolutely. To search out out new and authentic therapies, scientists square measure operating with ancient or people medications in parallel of recent medicine. *nigella sativa* has been used for healthful functions for hundreds of years. It originated from Southeastern Asia and also employed in ancient Egypt, Greece, geographic area and continent. In Islam, it's considered one among the best forms of healing medication out there. (Samajdar S, 2017) It's a flowering plant, of that seed is employed as a spice. The seed is termed black cumin in English, whereas in Latin it absolutely was known as 'Panacea' that means 'cure all'; in Arabic it's termed as 'Habbah Sawda' or 'Habbat el Baraka' translated as 'Seeds of blessing'. It's conjointly referred to as 'Kalo jeera' (in Bangladesh), 'Kalonji' (in India) and 'Hak Jung Chou' in (China). The seeds of NS are known for its carminative, stimulatory, protective, curative and diuretic activities and are generally employed in various recipes as spice and condiment due to its aroma particularly in Middle East, Asia, Italy, Germany and France Each seeds and oil extracted from this plant square measure employed in healthful functions. Kalonji (*Nigella sativa*) could be a dicotyledonous of ranunculacea is an incredible herb with a non secular background and usually grows within the Japanese Europe geographic area and Western Asia (Al-Bukhari and Sahi 1976). The seeds of *N. sativa* square measure the supply of the active ingredient of this plant. It is a black seed referred by Prophet mohammed as a panacea that's a remedy for all ailments however cannot stop ageing or death. The utilization of black seeds has been mentioned in numerous non secular and ethnic books. Black seeds square measure known because the curative black cumin within the holy Bible within the Unani-Tibb system

of drugs that originate from medical man, his modern anatomist and Ibn- sina has regarded black seed as a valuable remedy within the viscus and biological process disorder. Through thousands of years, till the time being, variant individuals within the Mediterranean region and much East countries use the oil of *N. sativa* seeds daily as a natural protecting and curative remedy. (Samajdar S, 2017).

## PLANT CHARACTERISTICS (Samajdar S, 2017)

### Taxonomic classification

Taxonomic classification	
Kingdom	Plantae
Subkingdom	Tracheobionta
Superdivision	Spermatophyta
Phylum	Magnoliophyta
Class	Magnoliopsida
Order	Ranunculales
Family	Ranunculaceae
Genus	<i>Nigella</i>
Species	<i>N. sativa</i>

### Common Names

Black cumin, Fennel Flower, Nutmeg Flower, Black seed, Black Caraway, Roman Coriander, Damascena, Devil in-the-bush, Wild Onion Seed, Kalonji, Kalojeera, Habat ul- Bakarrah.

### Habitat

*N. sativa* is native to Southern Europe, North Africa and Southwest Asia and it is cultivated in many countries in the world like Middle Eastern Mediterranean region, South Europe, India, Pakistan, Syria, Turkey, Saudi Arabia.

### Plant morphology

*N. sativa* is an annually flowering plant which grows from 25 cm to 95 cm in height, with finely divided leaf; the segments of the leaf are narrowly linear to threadlike. The flowers are delicate and are usually white, yellow, pink, pale blue or purple in colour. Number of petals present varies from 4-10. The fruit is large green n capsulated containing 3-8 united follicles, each having numerous black seeds.



**Fig: Nigella sativa plant**

**Characteristics of seed and flower:** (Samajdar S, 2017)

Macroscopically, seeds are small dicotyledonous, trigonus, angular, rugulose-tubercular, 2-3.5mm1-2 mm, black externally and white inside, odour slightly aromatic and taste bitter. Microscopically, transverse section of seed shows single layered epidermis consisting of elliptical, thick walled cells, covered externally by a papillosecuticle and filled with dark brown contents. Epidermis is followed by 2-4 layers of thick walled tangentially elongated parenchymatous cells, followed by a reddish brown pigmented layer composed of thick walled, rectangular elongated cells. Inner to the pigment layer, is present a layer composed of thick walled rectangular elongated or nearly columnar, elongated cells. Endosperm consists of thin walled, rectangular or polygonal cells mostly filled with oil globules. The powder microscopy of seed powder shows brownish black, parenchymatous cells and oil globules.



**Fig: Nigella Flower.**



**Fig: Nigella Seed**

**Chemical composition of black seeds** (Samajdar S, 2017)

Many active compounds have been isolated, identified and reported in different varieties of black seeds. The most important active compounds are thymoquinone (30%-48%), thymohydroquinone, dithymoquinone, p-cymene (7%-15%), carvacrol (6%-12%), 4-terpineol (2%-7%), t-anethol (1%-4%), sesquiterpene longifolene (1%-8%)  $\alpha$ -pinene and thymol etc. Black seeds also contain some other compounds in trace amounts. Seeds contain two different types of alkaloids; i.e. isoquinoline alkaloids e.g. nigellicimine and nigellicimineN-oxide and pyrazol alkaloids or indazole ring bearing alkaloids which include nigellidine and nigellicine.

Moreover, *N. sativa* seeds also contain alpha-hederin, a water soluble pentacyclic triterpene and saponin, a potential anticancer agent. Some other compounds e.g. carvone, limonene, citronellol were also found in trace amounts. Most of the pharmacological properties of *N. sativa* are mainly attributed to quinone constituents, of which TQ is the most abundant. On storage, TQ yields dithymoquinone and higher oligo condensation products. The seeds of *N. sativa* contain protein (26.7%), fat (28.5%), carbohydrates (24.9%), crude fibre (8.4%) and total ash (4.8 %). The seeds are also containing good amount of various vitamins and minerals like Cu, P, Zn and Fe etc. The seeds contain carotene which is converted by the liver to vitamin A. Root and shoot are reported to contain vanillic acid.

The seeds reported to contain a fatty oil rich in unsaturated fatty acids, mainly linoleic acid (50-60%), oleic acid (20%), eicodadienoic acid (3%) and dihomolinoleic acid (10%). Saturated fatty acids (palmitic, stearic acid) amount to about 30% or less.  $\alpha$ -sitosterol is a major sterol, which accounts for 44% and 54% of the total sterols in Tunisian and Iranian



varieties of black seed oils respectively, followed by stigmasterol (6.57-20.92% of total sterols).

Examples of various other reported chemical components includes nigellone, avenasterol-5-ene, avenasterol-7-ene, campesterol, cholesterol, citrostadienol, cycloeucaleanol, gramisterol, lophenol, obtusifoliol, stigmastanol, stigmasterol-7-ene,  $\beta$ -amyrin, butyrospermol, cycloartenol, 24-methylene-cycloartanol, taraxerol, tirucallol, 3-O- $\beta$ -D-xylopyranosyl(1 $\rightarrow$ 3)- $\alpha$ -L-rhamnopyranosyl(1 $\rightarrow$ 2)- $\alpha$ -L-arabino-pyranosyl]-28-O- $\alpha$ -L-rhamnopyranosyl (1 $\rightarrow$ 4)- $\beta$ -D-glucopyranosyl(1 $\rightarrow$ 6)- $\beta$ -D-glucopyranosyl] hederagenin, volatile oil (0.5-1.6%), fatty oil (35.6-41.6%), oleic acid, esters of unsaturated fatty acids with C15 and higher terpenoids, esters of dehydrostearic and linoleic acid, aliphatic alcohol,  $\beta$ -unsaturated hydroxy ketone, hederagenin glycoside, melanthin, melanthinigenin, bitter principle, tannin, resin, protein, reducing sugar, glycosidal saponin, 3-O- $\beta$ -D-xylopyranosyl-(1 $\rightarrow$ 2)- $\alpha$ -L-rhamnopyranosyl-(1 $\rightarrow$ 2)- $\beta$ -D-glucopyranosyl]-11-methoxy-16,23-dihydroxy-28-methyl-olean-12-enoate, stigma-5, 22-dien-3- $\beta$ -D-glucopyranoside, cycloart-23-methyl-7, 20, 22-triene-3 $\beta$ , 25-diol, nigellidine-4-O-sulfite, N. mines A3, A4, A5, C, N. mines A1, A2, B1, and B2.

#### **Traditional uses of folk remedies** (Samajdar S, 2017).

*N. sativa* has been traditionally used for the treatment of a variety of disorders, diseases and conditions pertaining to respiratory system, digestive tract, kidney and liver function, cardiovascular system and immune system support, as well as for general well-being. Avicenna refers to black seeds in the 'The Canon of Medicine', as seeds stimulate the body's energy and helps recovery from fatigue and dispiritedness. Black seeds and their oil have a long history of folklore usage in Indian and Arabian civilization as food and medicine. The seeds have been traditionally used in Southeast Asian and the Middle East countries for the treatment of several diseases and ailments including asthma, bronchitis, rheumatism and related inflammatory diseases. Its many uses have earned *Nigella* the Arabic approbation 'Habbatul barakah', meaning the seed of blessing. A tincture prepared from the seeds is useful in indigestion, loss of appetite, diarrhoea, dropsy, amenorrhoea and dysmenorrhoea and in the treatment of worms and skin eruptions. Externally the oil is used as an antiseptic and local anesthetic. Roasted black seeds are given internally to stop the vomiting.


## MATERIALS AND METHODS

### Collection

Seeds are collected from local market of Bhadrawati, District -Chandrapur, Vidarbha region of India.

### Extraction

The seeds of *Nigella sativa* are crushed to coarse powder using mortar pestle and passing through sieve. The extraction is done using four different solvents namely Petroleum Ether, Chloroform, Methanol, Hydro-alcoholic solution (20% DM Water in Methanol).



### Extraction process in flow diagram

#### Equipments used are.

1. Speed Extractor: Grounded Powder is loaded in the four chambers along with purified sand. The lead taking solvent is put in pure covered solvent. After three cycles of automatic extraction the equipment. The process continues for all four solvents. The extracts are stored in clean dried container.
2. Soxhlet Apparatus: Course the seeds of *Nigella sativa* and add with solvent in the chamber. After four cycles it completed. The extracts are stored in clean dried container.
3. Aspirator: Dried grounded powders are put in closed glass aspirator. Solvent in same order is poured over the powder so that all powder is in direct contact with solvent. The aspirator is kept overnight undisturbed with solvent. Next day the extract is collected and stored.

All the extracts are subjected to concentration using Rotavapor equipment until only concentrated oil or sticky solid is left. The hydro-alcoholic portion is fractionated against Petroleum ether, ethyl acetate and chloroform. The organic portion is collected and dried until minimum drop of oily substance is collected. They are mixed with parent extract and stored in cool and dry place. The only left water part is dried to form solid.

#### CHEMICAL TESTS: (Samajdar S, 2017)

Test Perform	Chemicals Test	Observation	Result
Test for Alkaloids	Saturated solution of picric acid+ filtrate	Yellow colour precipitate	Positive
Test for fixed oil	Drops of filtrate on Filter paper	Stain on filter paper	Positive
Test for volatile oil	1.Drops of filtrate on Filter paper	Permanent stain on filter paper	Positive
	2. Filtrate+ Sudan III dye	Red colour obtained	Positive
Test for Tannins	5% solution of FeCl <sub>3</sub> + filtrate	Green colour or yellow obtained	Positive
Test for Flavonoids	NaoH solution+ filtrate	Yellow colour obtained	Positive
Test for Glycoside	dilute H <sub>2</sub> SO <sub>4</sub> +filtrate + 5% solution of NaoH + add Fehiling solution (A+B)	Red colour produced	Positive
Test for Steroids	Conc. Sulphuric acid+ filtrate	Yellow colour on top	Positive
Test for Triterpenes	Conc. Sulphuric acid+ filtrate	Green colour on bottom produced	Positive

Result Tests	Pet. Ether Ext.	CHCl <sub>3</sub> Ext.	Methanolic Ext.	Water Ext.
Alkaloid Test	+	+	+	+
Fixed oil Test	+	+	+	-
Volatile oil Test	+	-	+	-
Test for tannin	-	-	-	-
Flavonoids Test	+	+	-	-
Glycoside test	-	-	-	-
Steroid test	+	+	+	+
Triterpenes test	+	-	+	+

#### CONCLUSION

From the above mentioned result we can conclude that all proportion of extract contains alkaloids, fixed oil, volatile oil, flavonoids stark absence of glycoside and tannins rest are present in some and absent in some.



**ACKNOWLEDGEMENT**

The authors are thankful to the Principal of Hi-Tech College of Pharmacy, Chandrapur (Maharashtra), India for providing necessary facilities to carry out this work. The authors are also thankful to the Principal of MVP College of Pharmacy, Nashik Maharashtra (India) for providing necessary facilities to carry out this work.

**REFERENCES**

1. Ali, M.A., Sayeed, M.A., Alam, M.S., Yeasmin, M.S., Khan, A.M., Muhamad, I.I, 2012. Characteristics of oils and nutrient contents of *Nigella sativa* Linn. and *Trigonella foenum-graecum* seeds. *Bull. Chem. Soc. Ethiop*, 26: 55–64.
2. Al-Bukhari, M. and A. Sahi (1976). "The collection of authentic sayings of Prophet Mohammad (peace be upon him), division 71 on medicine." Hilal Yayinlari, Ankara, Turkey.
3. Hedrick, U.P., Sturtevant, E.L., 1972. Sturtevant's Edible Plants of the World. Dover Publication Inc., New York, 388–389.
4. Khare, C. (2004). "Encyclopedia of Indian Medicinal Plants: Rational Western therapy, Ayurvedic and other Traditional usage." Springer, Germany, ISBN, 3: 540-20033.
5. Sharma, P., et al. (2000). Database on medicinal plants used in Ayurveda.
6. Riaz, M., Syed, M., Chaudhary, F.M., 1996. Chemistry of the medicinal plants of the genus *Nigella*. *Hamdard Medicus*, 39: 40–45.
7. Samajdar S; Extraction and chemical tests on *Nigella Sativa* l. collected from Punjab region of India; *Pharma Tutor*, 2017; 5(12): 54-57.
8. Nimet KARA, Duran KATAR, Hasan BAYDAR; Yield and quality of black cumin (*nigella sativa* l.) Populations: The effect of ecological conditions; *turk J Field Crops*, 2015; 20(1): 9-14.