

QUALITATIVE AND QUANTITATIVE ESTIMATIONS OF PHYTOCONSTITUENTS FROM *PHOENIX DACTYLIFERA* ROOTS

¹*Shakeeb Akhtar Nehal Ahmad, ²Dr. Rakesh Kumar Jat and ³Dr. Rashid Akhtar

¹Research scholar, JJT University, Vidyanagari, Churela, Jhunjhunu, Rajasthan 333001.

²Principal Department of Pharmacy, JJT University, Vidyanagari, Churela, Jhunjhunu, Rajasthan 333001.

³Associate professor, Department of Pharmacognosy, Royal college of Pharmaceutical Education and Research, Sayne Khurd, , Malegaon, Dist. Nashik, Maharashtra.

Article Received on
28 Oct. 2019,

Revised on 18 Nov. 2019,
Accepted on 08 Dec. 2019,

DOI: 10.20959/wjpr20201-16506

*Corresponding Author

**Shakeeb Akhtar Nehal
Ahmad**

Research Scholar, JJT
University, Vidyanagari,
Churela, Jhunjhunu,
Rajasthan 333001.

ABSTRACT

The date palm (*Phoenix dactylifera*) has been known and used for medicinal purposes since ancient times. The edible part of the date palm tree has been recognized to possess many medicinal properties when consumed alone or in mixture with other medicinal herbs. In recent years, a huge interest in the abundant health promoting properties of date fruits had led to many pharmacological studies (in-vitro and in-vivo) as well as the identification and quantification of different classes of phytochemicals. The root part was selected. Morphological characteristic were performed and transverse section was taken and observed under microscope. Extraction was done by using methanol as solvent in Soxhlet apparatus. Extract was subjected

for different phytochemical constituent. Total flavonide and terpenoid were determined and it was observed that Phoenix dactylifera root contains 7.25% total flavonide and 0.25% terpenoide.

KEYWORD: *Phoenix dactylifera* root, Methanolic extract, Flavonide, terpenoid.

INTRODUCTION

Many plants, including some of the commonly consumed herbs and spices in our daily food, can be safely and effectively used to prevent and/or treat some health concerns. Foreexample, caffeine the active ingredient found in coffee beans (*Coffea*), shows biological activity in the

treatment of the central nervous system (CNS) disorders indole-3-carbinol, and 3,3'-diindolylmethane are both broccoli (*Brassica oleracea*) derived phytochemicals with potential anti-cancer activity, and resveratrol, isolated from grape (*Vitisvinifera*), is reported to extend lifespan and provide cardio-neuro-protective, anti-diabetic, and anti-cancer effects. Date palm fruits possess high nutritional and therapeutic value with significant antioxidant, antibacterial, antifungal, and anti-proliferative properties.^[1] Natural products can be good remedies because they are inexpensive and easy to access.^[2]

The date palm (*Phoenix dactylifera*) has been known and used for medicinal purposes since ancient times.^[4] Date palm tree (*Phoenix dactylifera* L.) is considered as one of the oldest and main staple and ancient crops in Southwest Asia and North Africa. Besides, dates can be grown in Australia, Mexico, South America, southern Africa, and the United States, especially in southern California, Arizona, and Texas. Date palm tree belongs to Arecaceae family (Angiosperms, monocotyledon) consisting of about 200 genera and more than 2,500 species. Phoenix (Coryphoideaaphoeniceae) is one of the genera with approximately 14 species which are native to the tropical or subtropical regions of southern Asia or Africa, including *Phoenix dactylifera* L. The name of the species *dactylifera* means “finger-bearing” which refers to the fruit clusters produced by this plant. *Dactylifera* is a grouping of the Greek word *dactylus*, means “finger,” and the Latin word *ferous*, mean “bearing”. Very recently the whole genome of date palm tree was re-sequenced yields insights into diversification of a fruit tree crop.^[1] Dates are implicated to possess medicinal properties in addition to its nutritional value. Several studies have reported date fruit with a wide range of bioactivities, such as antioxidant activity due to the presence of phenolics, carotenoids and anthocyanins, antimutagenic, anti-inflammatory, antihyperlipidemic, antibacterial and antifungal activities. Date fruit consist of 70% carbohydrates, most of which is in the form of sugars. Because of this the fruits are a high source of energy and it is approximated that 100 g of the flesh can provide 314 kcal of energy. Drying of date decreases the water activity and this increases the sugar concentration. Because of this the shelf life of dry dates are high and are available for extended periods of time. The fruits are also used as a sweetener in the preparation of beer. The date pits which are a waste product have been used for centuries in the Arab world to make caffeine-free drink. Recently, date pit powders are also marketed and are a source of choice to people preferring a non-caffeinated coffee with coffee-related flavor. Date pits are also ground and added to the feed of domesticated animals and observations suggest they are devoid of any harmful effects. Date pits are rich in protein (5.1 g/100 g), fat

(9.0 g/100 g), dietary fiber (73.1 g/100 g), phenolics (3942 mg/100 g), and antioxidants (80,400 $\mu\text{mol}/100\text{ g}$), and may be of use in enhancing the nutritional value of incorporated food products. The most important quality attributes to grade dates are color, flavor (sugar level), moisture (26–30%) and absence of defects such as insect, damage, cracks and surface damage. Date fruit is good source of high nutritional value food. In more details, carbohydrates forms 70% of date fruit and are mostly fructose and glucose in equal ratio while date proteins are rich in amino acids that contain acidic side chain but poor in methionine and cysteine, which their side chain composed of sulfur. Minerals in date fruits are calcium, iron, magnesium, selenium, copper, phosphorus, potassium, zinc, sulfur, cobalt, fluorine, manganese, and boron. The edible part of the date palm tree has been recognized to possess many medicinal properties when consumed alone or in mixture with other medicinal herbs. In recent years, a huge interest in the abundant health promoting properties of date fruits had led to many pharmacological studies (in-vitro and in-vivo) as well as the identification and quantification of different classes of phytochemicals. The date fruits are highly nourishing and may have numerous potential health benefits. The protective effects of fruits against chronic diseases are ascribed to bioactive non-nutrients called phytochemicals. Phytochemicals have gained increased interest among several investigators, including clinicians due to their antioxidant activity, cholesterol-lowering properties, and other potential health benefits such as chemoprevention of cancer, prevention of diabetes and cardiovascular diseases.^[3] Numerous beneficial health effects have long been associated with date fruit, including antioxidant, anti-mutagenic and anti-inflammatory activity, and protection of the gastric mucosa against damaging effects of stomach acid.^[1] Hepatoprotective activity has also been linked to date fruit, including reduced alkaline phosphatase levels. These effects have been linked to the presence of anthocyanins, ferulic acid, caffeic acid, quercetin and proanthocyanidins.^[6] Every part of the date-palm has good use: The wood and leaves provide timber and fabric for houses and fences; the leaves are also used for making ropes, baskets etc; the stalk is used as fuel. The fruits famous for their delicious and sweet taste are taken directly or processed to produce vinegar, pickle, bakery items and flavours. The date seeds are used as cattle feed after they have been softened by soaking and crushing. The date-palms are usually un branched. It is interesting to note that the branching occurs only in the male plants! There are typically a dozen bunches of dates per tree. A bunch weighs about seven kilograms and has a thousand dates. The life of a date tree is over hundred and fifty years. There are about a thousand varieties of dates. Some of these are facing possible extinction. The date fruits vary in size, shape and colour. Each of these

numerous varieties have different name in Arabic. According to the estimates of the World Food Agricultural Organization, there are over hundred million date-palms in the world, producing two million tons of dates each year. About 65% of these are grown in the Arab countries. Let us have a closer look at this familiar fruit, which is also known as the *tree of life* and *king of the oasis*. The botanical name of the date-palm is *Phoenix dactylifera* Linn. In Arabic the date-palm is known as *Nakhl*, and the fruit is known as *Tamar*. We shall note the names in several other languages: *Tamar* (Hebrew), *Khajur* (Hindi, Urdu, and several Indian languages), *Khurma* (Persian, Urdu), *Kharjur* (Sanskrit), *Khejur* (Bengali), *Finik* (Russian), *Datil* (Spanish), *Date* (Italian), *Datteir* (French), *Daten* (German), *Datum* (Dutch), *Datas* (Portugese) and so on.^[4]

The *Phoenix dactylifera* consists of polyphenols like cinnamic acid and their derivatives in which coumaric acid, ferulic acid, and sinapic acids are major compounds. The darker color variety of the date fruit has a higher concentration of polyphenols compared to the lighter variety. The date palm kernel contains phytohormones; thus, it produces a significant anti-wrinkle effect and can be used in antiaging products. Date fruits are rich in phenolic and also possess antioxidant activity. Date seeds can also be used in skin products, as they can yield moisturizing oil with different essential fatty acids.^[4]

MATERIALS AND METHODS

Plant Material

Root of *Phoenix dactylifera* was collected from Chandwad (District Nashik, M.S.) and authenticated from the department of botany, Dr. Babasaheb Ambedkar University, Aurangabad.

Morphology

The external morphology was studied according to reported methods. Macroscopic study was performed by using simple microscope. The colour, odour, taste, size and shape of flowers and roots were determined.

Transverse section & powder characteristics

The free-hand section of root were prepared from fresh plant material and finally stained with various staining reagents as per standard procedures. The disaggregation of plant material was performed by reported method. In brief, scales were disaggregated by means of boiling in an aqueous solution of NaOH (5% w/v) for 5 min. After cooling and washing with water,

pieces were treated with an aqueous solution of chromic acid (25 % v/v) for 30 min at room temperature. The section was cleared with chloral hydrate solution, stained with phloroglucinol-hydrochloric acid (1:1) and toluidine blue.^[26] Powdered drug was used for the observation of powder microscopical character. The powdered drug was separately treated with phloroglucinol-hydrochloric acid(1:1)solution, acetic acid and iodine solution to determine the presence of the lignified fibres, Calcium oxalate crystals and starch grains respectively. A series of digital images captured using a Motic Digital microscope fitted with 1/3" CCD camera imaging accessory and using Motic Images 2000 (1.3 version) analysis software. The micrometric data were generated from average of 30 measurements for each sample and expressed as lower limit mean+SD.^[8]

Fluorescence analysis

Fluorescence study of powdered drug of root of *Phoenix dactylifera* was performed by following a standard procedure.^[28] In this study the powdered root was treated with various acidic and basic solvents and observed in UV visible chamber under short and long wavelength region simultaneously.^[24, 25] Fluorescence is an important factor which reveals various chemical constituents show fluorescence in the visible range in day light. UV light gives fluorescence to many natural products such as berberin (an alkaloid) which do not generally produce fluorescence in day light. If the active constituents are not themselves fluorescent, they can be converted into fluorescent derivatives by using reagents. Hence it plays an important role for the pharmacognostical parameter for the evaluation of crude drugs.^[29]

Extraction of root of *Phoenix dactylifera*

50 g of powdered root of *Phoenix dactylifera* was packed in white cotton bags and then extracted successively using the following solvents in a Soxhlet apparatus. The solvents used for extraction were petroleum ether (60-80 °C), chloroform and methanol.

Each time before extracting with the next solvent, the powdered material was dried in an air oven below 50 °C. The extractive values and preliminary phytochemical studies were carried out on these successive extracts.

Qualitative estimation

In the present study, the qualitative chemical tests were performed on different successive extracts of *Phoenix dactylifera*. For the phytoconstituents such as alkaloids, glycosides,

tannins, resins, carbohydrates, etc. The results obtained on examination of successive extracts are indicated in the following table.^{[36] [37]}

Quantitative Estimations

• Estimation Of Total Phenolic Content

To determine total phenolic content from the methanolic extract of root of *Phoenix dactylifera*, calibration curve of standard gallic acid of 20, 40, 60, 80 and 100 mg/ml was prepared in water and mg/ml of methnolic extract of root *Phoenix dactylifera* was prepared simultaneously. Each sample was mixed with 0.25 ml of Folinicocalteu reagent and 1.25 ml sodium carbonate solution. The mixtures were allowed to react for 40 minutes at room temperature. After the reaction period the blue color ease measured at 725 nm on UV visible spectrophotometer of LABINDIA 3000+ and calculated the amount of total phenolic content from calibration curve as gallic acid equivalent by using following formula.

$$T=C.V/M$$

Where T= Total content of phenolic compound in mg per gram of plant extract.

C= Concentration of gallic acid established from the calibration curve mg/ml.

V=Volume of extract in milliliter

M= gram weight of plant extract.^[38, 39]

• Estimation Of Total Flavonoid Content

An aliquot (1 ml) of standard solution of quercetin (20, 40, 60, 80,100 µg/ml) was added to 10 ml volumetric flask containing 4 ml of 5% NaNO₂ into it. After 5 minute 0.3 ml of 10% AlCl₃ was added. Then 2 ml of 1 M NaOH was added and the total volume was made up to 10 ml with distilled water. Same dilutions were also prepared for the test solution. Blank determination was done by using methanol in place of test or standard solutions.

Mixed well and taken the absorbance at 358 nm against blank. From the obtained standard curve of quercetin the total flavonoids content of methanolic extract of root of *Phoenix dactylifera* was determined.

RESULTS AND DISCUSSION

Morphology

External Morphology was studied by according to reported methods. The obtained results are as follows.




Figure 01: Roots of *Phoenix dactylifera*.

Table 01: Result of Morphological study of *phoenix dactylifera* root.

Test Parameters	Observation
Colour	Whitish Brown
Odour	Characteristic
Taste	Sweet
Shape	Long irregular

Transverse section of root of *Phoenix dactylifera*

The microscopy of root of *Phoenix dactylifera* was performed by using monochromatic staining agent like Phloroglucinol and HCL in proportion of 1:1. It revealed presence of typical root characteristics as central pith, few layers of cork cell followed by cortex. Lignified vascular tissues were also observed. In epidermis few layers of cork is observed and in endodermis cortex, vascular tissue, and pith were observed.



Cr= Cortex, PH= Phloem, ED= Endodermis, VT= Vascular tissue

Figure 02: Transvers section of *Phoenix dactylifera* root.

Fluorescence analysis

Present fluorescence analysis on powdered root of *Phoenix dactylifera* Shows characteristic colors with reagents like alcohol, 10% NaOH, and 10% HNO₃, and 50% H₂SO₄. The result is depicted in table no 2.

Table 02: Fluorescence analysis of root.

Fluorescence study		
Reagents	Long UV light	Short UV light
Powder as such	Yellow	Yellowish green
Powder in distilled water	Yellow	Yellowish green
Powder in Absolute alcohol	Pale green	green
Powder In 10% NaOH	Green	Dark green
Powder in 10% HNO ₃	Light green	green
Powder in 50% H ₂ SO ₄	Yellow	Pale yellow

Qualitative estimation of phytoconstituent**Table 03: Result of Phytochemical tests of extracts.**

Sr.No	Constituents	Tests	Pet. Ether	Chloroform	Methanol
1.	ALKALOIDS	Mayer's test	-	-	-
		Dragordraff's test	-	-	-
		Hager's test	-	-	-
		Wagner's test	-	-	-
2.	STEROLS	Liebermann'sBurchard test	-	-	+
		Salkowski's -	-	+	+
3.	CARBOHYDRATES AND GLYCOISDES	Molisch's test	-	+	+
		Fehling's test	-	+	+
		Benedict's test	-	+	+
		Borntrager's test	-	+	+
4.	FIXED OILS AND FATS	Spot test	-	-	-
		Saponification test	-	-	-
5.	PHENOLIC COMPOUND	FeCl ₃ test	--	-	+
6.	PROTEIN AND AMINOACIDS	Biuret test	-	-	-
		Ninhydrin test	-	-	-
		Xanthoprotein test	-	-	-
		Millon's test	-	-	-
7.	TRITERPINOID AND SAPONINS	Foam test	-	-	+
		Haemolysis test	-	-	+
8.	TANNINS	Gelatin test	-	-	-
		FeCl ₃ test	-	-	-
9.	GUMS AND MUCILAGE	Precipitation with 90% alcohol	-	-	-
10.	FLAVONOIDS	Aqueous NaOH	+	+	+
		Conc. H ₂ SO ₄	+	+	+

Quantitative estimation

- **Total Phenolic content**

For quantitative estimations, total phenolic content, total flavonoid content were determined from methanolic extract of root of *Phoenix dactylifera*.

Each quantitative estimation was done by using standard samples. Results of each quantitative parameter are given below.

Table 04: Absorbance observed in estimation of Total Phenolic content from root.

Sr. No.	Standard (Gallic acid mg/ml)	Absorbance			Mean
		I	II	III	
1	20	0.275	0.273	0.278	0.275
2	40	0.56	0.56	0.57	0.563
3	60	0.887	0.888	0.884	0.886
4	80	1.188	1.190	1.185	1.187
5	100	1.458	1.460	1.457	1.458
RPD	Unknown	1.042	1.042	1.043	1.042

(RPD = Root of Phoenix dactylifera)




Figure 03: Calibration curve of standard Gallic acid.

Determination of total concentration of Phenolic compound content was calculated on the basis of standard calibration curve of Gallic acid. Total Phenolic content from root of PD was calculated as 7.10%.

- Total flavonoid content

Table 05: Absorbance observed in estimation of Total flavonoid content from root PD.

Sr. No.	Gallic acid ($\mu\text{g/mL}$)	Absorbance			Mean
		I	II	III	
1	20	0.053	0.055	0.054	0.054
2	40	0.133	0.132	0.133	0.132
3	60	0.223	0.226	0.224	0.224
4	80	0.309	0.311	0.308	0.309
5	100	0.377	0.374	0.379	0.376
RPD	Unknown	0.282	0.282	0.281	0.281




Figure 04: Calibration curve of standard Quercetin.

CONCLUSION

Natural products can be good remedies because they are inexpensive and easy to access. Date palms possess high nutritional and therapeutic value with significant antioxidant, antibacterial, antifungal, and anti-proliferative properties. In above study root part of *Phoenix dactylifera* was selected because very rear work has been done on root. Methanolic extract was subjected for phytoconstituent and it was observe that the root contains flavonoid and terpenoid.

REFERENCES

1. Reem A., Jawhara H., Jawaher S. M., Al-Nadabi, Badria I. Al-Shihi, Younis B., Date palm tree (*Phoenix dactylifera*) natural product and therapeutic options. *Frontiers in plant science*, 2017; 8: 845.
2. Ali Hafez, Hazem M. S., Mohamed M., Soad K. Al Jaouni. Date Palm (*Phoenix dactylifera*) Protection and Remedy Food. *Journal of Nutraceuticals and Food Science*, 2016; 29: 01-10.

3. Manjeshwar S.B., Bantwal R. B., Shaun M.k., Harshith P. B., Praveen V. A review of the chemistry and pharmacology of the date fruits (*Phoenix dactylifera* L.). *Food Research International*, 2016; 44: 112-122.
4. Hajira Khan, Sameen Ahmed Khan. Date palm revisited. *Research journal of pharmaceutical, biological and chemical*, 2016; 09: 10-20.
5. Saada M. Al-Orfi, Mona H. M., Norah Al- Atwai1, Huda Al Zaidi, Asma D., SumayyahD. Review nutrition properties and benefits of the date fruits. *Bulletin of the National Nutrition Institute of the Arab Republic*, 2012; 39: 97.
6. Emilia Alfaro-Viquez, Brent F. Roling, Christian G. Krueger, Charlene J. Rainey, Jess D. An extract from date palm fruit (*Phoenix dactylifera*) acts as co agonist ligand for the nuclear receptor FXR and differentially modulates FXR targets gene expression in vitro. *Plos one*, 2018; 13: 1-23.
7. Chuan-Rui Z., Saleh A. Aldosari , Polana S.P.V., Paraj S., Muraleedharan G. Nair. Health benefits of date fruit produced in Saudi Arabia based on in vitro anti-oxidant, anti-inflammatory and human tumor cell proliferation inhibitory assays. *Journal of the Saudi Society of Agricultural Sciences*, 2017; 16: 287-293.
8. Edwin S., Joshi S.B., Jain D.C. Comparative pharmacognostic studies on root powder of *Plumbagozeylanica* and *Plumbagorosea*. *Indian J. Nat Prod*, 2008; 24(2): 27–29.
9. Afzal k., Tarique Jamal K., Guthaman, Peter N.P., Adeel Chaudhary, Taha Kumosani, Mohammed Al-Qahtani. Anticancer effect of ajwa dates. *BMC Complementary and Alternative Medicine*, 2017; 17: 418.
10. Wan Iryani, Wan Ismail, Mohd Naim, Fadhli Mohd Radzi. Evaluation of benefits of date palm to the brain. *Alternative and integrative medicine*, 2013; 2357: 1-3.
11. Ranjitha Dhevi V. Sundar, Gayathri Segaran, Saranya S., Sugashini Settu, Lokesh Ravi. Bioactivity of *Phoenix dactylifera* seed and its phytochemical analysis. *International journal of green pharmacy*, 2017; 11: 292.
12. Sahar Rahmani, Hojjatullah, Parham R., Zahra S. Effect of hydro alcoholic *Phoenix dactylifera* extract on sleep and EEG in rat. *Avicenna journal of phytomedicines*, 2016; 6: 511-518.
13. Naskar S., Islam A. Shah P., Haldar P.K., Gupta M. In vitro in vivo anti oxidnt potential of hydromethanolic extract of *Phoenix dactylifera* fruit. *Journal of scientific research*, 2010; 2(1): 144-157.
14. Najla Bentrada, Rabéa G., Yamina Benmalek, Fatma Rahmania. Studies chemical composition and antimicrobial activities of bioactive molecules from date palm (*Phoenix*

- dactylifera) pollens and seeds. *AfrJ Tradit Complement Altern Med*, 2017; 14(3): 242-256.
15. Ali Ramazani, Marzieh Mashhadi, Mohammad Kamalinejad, Mahdi Tavakolizadeh, Fatemeh Balaghi, Maryam Nobarani, Mohammad Reza Eskandari, Farshad H. S. Phoenix dactylifera fruit induce cell proliferation A2780, A172 AND HFFF2 Cell lines. *Iranian journal of pharmaceutical sciences*, 2016; 12: 35-44.
 16. Mehrdad Shakiba, Ashraf Kariminik, Paria Parsia. Antimicrobial Activity of Different Parts of *Phoenix dactylifera*. *International journal of molecular and clinical biology*, 2011; 1: 107-111.
 17. Eva Agustina, Risa Purnamasari, Nova Lusiana. Effect of extract meat date to oxytocine level in pregnant mice (*Mus musculus L.*). *Humanistic Network for Science and Technology*, 2017; 2: 1-09.
 18. Soad Al-daihan, Ramesa Shafi Bhat. Antimicrobial activities of extract of leaf, fruit, seed and bark of *Phoenix dactylifera*. *African journal of biotechnology*, 2012; 11: 10021-10025.
 19. Rahmani, Salah Ali Y., Sauda M., Habeeb Ali, Arshad H. S., Amjad A. Khan. A review article on therapeutic effect of date fruit in the prevention of disease via modulation of anti-inflammatory, anti-oxidant and anti-tumor activity. *International Journal of clinical experiment and medicine*, 2014; 7(3): 483-491.
 20. Ibrahim A., Maged E., Ahmadk.k. Date Palm (phoenix dactylifera) fruit as a potential cardio protective agent. *Frontiers in pharmacology*, 2017; 8: 592.
 21. Shalam Mohamed Hussain, Exploring the possible role of date palm (phoenix dactylifera) extract in amelioration of stroke in rats. *Asian journal of Biomedical and Pharmaceutical science*, 2015; 5(51): 31-35.
 22. Shadab A., Rafeeq Alam K., Subia Jamil, Syeda A. Antidiabetic effect of native date fruit (phoenix dactylifera) normal and hyperglycemic rat. *Pak journal of pharmaceutical science*, 2017; 30(5): 1797-1802.
 23. Chase C.R., Pratt R.J. Fluorescence of powdered vegetable drugs. *Indian Journal of Experimental Biology*, 1949; 33(6): 428-432.
 24. Ansari S.H. *Essential of Pharmacognosy*. 1st edition, New Delhi, Birla publications Pvt. Ltd, 2006; 43-46.
 25. Sulaiman Al-Zuhair, Ali Dowaidar, Hassan Kamal. Inhibitory effect of dates extract on α amylase and α glycosidase enzyme relevant to non-insulin dependent diabetes mellitus. *Journal of biochemical tec*, 2010; 2(2): 158-160.

26. Seyyed Ali M., KowtharJ., Mohammadkazem G. Evaluation of the antidiabetic and antilipaemic activities of hydroalcoholic extract of Phoenix dactylifera palm leaves and its function on diabetic rats. Malaysian journal of medical science, 2010; 17(4): 4-13.
27. Fazal K., Tariq Jamal k., Gauthaman K., Peter Natesan Pushparaj, Adeel Chaudhary. Anti-cancer activity of ajwa dates. Complementry and alterntive medicines, 2017; 17: 2-10.
28. Khandelwal K.R. Practical Pharmacognosy. Nirali Prakashan, Pune, 2005; 149-153.
29. Mobsher A., IqraMasood, HafeezIqram. Pharmacological investigation of Phoenic dactylifera in azithromycin induced toxicity. International journal of Pharmacology, 2018; 14(1): 61-67.
30. Priyanka Verma. Effect of date in lowering LDL in geriatric cases. Health science journal, 2016; 5(8): 1-9.